

## Comprehensive compilation of Randomized Controlled Trials (RCTs) involving Psychodynamic treatments (PDT)

Peter Lilliengren, PhD, assistant professor  
Department of Psychology, Stockholm University, SWEDEN  
<http://w3.psychology.su.se/staff/pe/i/indexeng.html>

Last updated: 2016-08-17

#	Reference (primary and secondary articles)	Patient sample	PDT-model	Comparison group(s)
1.	<b>Abbass, A., Sheldon, A., Gyra, J., &amp; Kalpin, A.</b> (2008). Intensive short-term dynamic psychotherapy for DSM-IV personality disorders: a randomized controlled trial. <i>J Nerv Ment Dis</i> , 196, 211-216. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18340256">http://www.ncbi.nlm.nih.gov/pubmed/18340256</a>	<b>Personality disorders</b> (mixed)	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Davanloo, 1990)	<b>Minimal contact</b>
2.	<b>Ajlchi, B., Nejati, V., Town, J. M., Wilson, R., &amp; Abbass, A.</b> (2016). Effects of Intensive Short-Term Dynamic Psychotherapy on depressive symptoms and executive functioning in Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 204(7), 500-5. <a href="http://doi.org/10.1097/NMD.0000000000000518">http://doi.org/10.1097/NMD.0000000000000518</a>	<b>Depression</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Davanloo, 1990)	<b>Waitlist</b>
3.	<b>Alstrom, J. E., Nordlund, C. L., Persson, G., Harding, M., &amp; Ljungqvist, C.</b> (1984). Effects of four treatment methods on social phobic patients not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 97-110. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6148843">http://www.ncbi.nlm.nih.gov/pubmed/6148843</a>	<b>Social anxiety disorder (SAD)</b>	<b>Supportive dynamic psychotherapy</b> (Dewald, 1964)	<b>Prolonged exposure</b> (in vivo w. therapist)/ + <b>relaxation therapy</b>
4.	<b>Alstrom, J. E., Nordlund, G., Persson, G., Harding, M., &amp; Lundquist, C.</b> (1984). Effects of four treatment methods on agoraphobic women not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 1-17. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6147066">http://www.ncbi.nlm.nih.gov/pubmed/6147066</a>	<b>Agoraphobia</b> (in women not suitable for insight-oriented psychotherapy)	<b>Supportive dynamic psychotherapy</b> (Dewald, 1964)	<b>Prolonged exposure</b> (in vivo w. therapist)/ + <b>relaxation therapy</b>
5.	<b>Amianto, F., Ferrero, A., Pierò, A., Cairo, E., Rocca, G., Simonelli, B., ... Fassino, S.</b> (2011). Supervised team management, with or without structured psychotherapy, in heavy users of a mental health service with borderline personality disorder: a two-year follow-up preliminary randomized study. <i>BMC psychiatry</i> , 11(1), 181. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260092&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260092&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Borderline personality disorder</b>	<b>Sequential Brief Adlerian Psychodynamic Psychotherapy</b> (SB-APP; Fassino, 2008; 2009)	<b>Supervised team management</b>
6.	<b>Andersson, G., Paxling, B., Roch-Norlund, P., Ostman, G., Norgren, A., Almlöv, J., ... Silverberg, F.</b> (2012). Internet-Based Psychodynamic versus Cognitive Behavioral Guided Self-Help for Generalized Anxiety Disorder: A Randomized Controlled Trial. <i>Psychotherapy and psychosomatics</i> , 81(6), 344-355.	<b>Generalized Anxiety Disorder (GAD)</b>	<b>IPDT</b> (SUBGAP; Silverberg, 2005)	<b>ICBT</b> (Paxling, 2011)

	<a href="http://www.ncbi.nlm.nih.gov/pubmed/22964540">http://www.ncbi.nlm.nih.gov/pubmed/22964540</a>			
7.	<p><b>Arnevik, E., Wilberg, T., Urnes, O., Johansen, M., Monsen, J. T., &amp; Karterud, S. (2009).</b> Psychotherapy for personality disorders: short-term day hospital psychotherapy versus outpatient individual therapy - a randomized controlled study. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 24(2), 71–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19097870">http://www.ncbi.nlm.nih.gov/pubmed/19097870</a></p> <p>Arnevik, E., Wilberg, T., Urnes, Ø., Johansen, M., Monsen, J. T., &amp; Karterud, S. (2010). Psychotherapy for Personality Disorders: 18 Months' Follow-Up of the Ullevål Personality Project. <i>Journal of Personality Disorders</i>, 24(2), 188–203. <a href="http://guilfordjournals.com/doi/pdf/10.1521/pepi.2010.24.2.188">http://guilfordjournals.com/doi/pdf/10.1521/pepi.2010.24.2.188</a></p>	<b>Personality disorders (mixed)</b>	<b>Short-Term Day Hospital Psychotherapy (DHP; no specific reference)</b>	<b>Outpatient Individual Psychotherapy</b>
8.	<p><b>Bachar, E., Latzer, Y., Kreitler, S., &amp; Berry, E. M. (1999).</b> Empirical comparison of two psychological therapies. Self psychology and cognitive orientation in the treatment of anorexia and bulimia. <i>J Psychother Pract Res</i>, 8, 115-128. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10079459">http://www.ncbi.nlm.nih.gov/pubmed/10079459</a></p>	<b>Bulimia</b>	<b>Self Psychological Treatment (based on Kohut, 1984)</b>	<b>Cognitive Orientation Treatment (Kreitler &amp; Kreitler, 1990)</b>
9.	<p><b>Baldoni, F., Baldaro, B., &amp; Trombini, G. (1995).</b> Psychotherapeutic perspectives in urethral syndrome. <i>Stress Medicine</i>, 11(July 1994), 79–84. <a href="http://onlinelibrary.wiley.com/doi/10.1002/smi.2460110115/abstract">http://onlinelibrary.wiley.com/doi/10.1002/smi.2460110115/abstract</a></p>	<b>Urethral syndrome</b>	<b>STDP (Malan, 1979; Davanloo, 1980)</b>	<b>TAU</b>
10.	<p><b>Balottin, U., Ferri, M., Racca, M., Rossi, M., Rossi, G., Beghi, E., ... Termine, C. (2014).</b> Psychotherapy versus usual care in pediatric migraine and tension-type headache: a single-blind controlled pilot study. <i>Italian Journal of Pediatrics</i>, 40(1), 6. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24444140">http://www.ncbi.nlm.nih.gov/pubmed/24444140</a></p>	<b>Idiopathic headache (children/adolescents 6-18 yrs)</b>	<b>STPP (Aliprandi et al, 1990)</b>	<b>TAU</b>
11.	<p><b>Barber, J. P., Barrett, M. S., Gallop, R., Rynn, M. a, &amp; Rickels, K. (2011).</b> Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. <i>The Journal of clinical psychiatry</i>, 1–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22152401">http://www.ncbi.nlm.nih.gov/pubmed/22152401</a></p> <p>Zilcha-Mano, S., Dinger, U., McCarthy, K. S., Barrett, M. S., &amp; Barber, J. P. (2013). Changes in well-being and quality of life in a randomized trial comparing dynamic psychotherapy and pharmacotherapy for major depressive disorder. <i>Journal of affective disorders</i>, 1–5. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24176534">http://www.ncbi.nlm.nih.gov/pubmed/24176534</a></p>	<b>Depression</b>	<b>Supportive Expressive Therapy (Luborsky, 1984)</b>	<b>1. Placebo</b> <b>2. SSRI/SNRI</b>
12.	<p><b>Barkham, M., Rees, A., Shapiro, D. A., Stiles, W. B., Agnew, R. M., &amp; Halstead, J., et al. (1996).</b> Outcomes of time-limited psychotherapy in applied settings: replicating</p>	<b>Depression</b>	<b>Psychodynamic-interpersonal Therapy (Hobson, 1985)</b>	<b>CBT (no reference)</b>

	the Second Sheffield Psychotherapy Project. <i>J Consult Clin Psychol</i> , 64, 1079-1085. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8916639">http://www.ncbi.nlm.nih.gov/pubmed/8916639</a>			
13.	<b>Barkham, M., Shapiro, D. A., Hardy, G. E., &amp; Rees, A. (1999).</b> Psychotherapy in two-plus-one sessions: outcomes of a randomized controlled trial of cognitive-behavioral and psychodynamic-interpersonal therapy for subsyndromal depression. <i>J Consult Clin Psychol</i> , 67, 201-211. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10224730">http://www.ncbi.nlm.nih.gov/pubmed/10224730</a>	<b>Depression</b> (subsyndromal)	<b>Psychodynamic-interpersonal Therapy</b> (3 sessions; Barkham & Shapiro, based on Hobson, 1985)	<b>CBT</b> (Beck et al 1979; Beck, 1995)
14.	<b>Bassett, D. L., &amp; Pilowsky, I. (1985).</b> A study of brief psychotherapy for chronic pain. <i>J Psychosom Res</i> , 29, 259-264. <a href="http://www.sciencedirect.com/science/article/pii/0022399885900522">http://www.sciencedirect.com/science/article/pii/0022399885900522</a>	<b>Chronic pain</b>	<b>12 sessions PDT</b> (no reference)	<b>Supportive psychotherapy</b> (no reference)
15.	<b>Bastos, A. G., Guimaraes, L. S. P., &amp; Trentini, C. M. (2014).</b> The efficacy of long-term psychodynamic psychotherapy, fluoxetine and their combination in the outpatient treatment of depression. <i>Psychotherapy Research : Journal of the Society for Psychotherapy Research</i> , (July 2014), 1–13. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25041333">http://www.ncbi.nlm.nih.gov/pubmed/25041333</a>  <b>Bastos, A. G., Pinto Guimarães, L. S., &amp; Trentini, C. M. (2013).</b> Neurocognitive changes in depressed patients in psychodynamic psychotherapy, therapy with fluoxetine and combination therapy. <i>Journal of Affective Disorders</i> , 1–10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24103853">http://www.ncbi.nlm.nih.gov/pubmed/24103853</a>	<b>Depression</b>	<b>Long-term psychodynamic psychotherapy (LTPP;</b> Gabbard, 2004; 2010)	<b>1. Fluoxetine alone</b>  <b>2. LTPP + fluoxetine</b>
16.	<b>Bateman, A., &amp; Fonagy, P. (1999).</b> Effectiveness of partial hospitalization in the treatment of borderline personality disorder: a randomized controlled trial. <i>The American Journal of Psychiatry</i> , 156(10), 1563–9. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10518167">http://www.ncbi.nlm.nih.gov/pubmed/10518167</a>  <b>Bateman, A., &amp; Fonagy, P. (2008).</b> 8-Year Follow-Up of Patients Treated for Borderline Personality Disorder: Mentalization-Based Treatment Versus Treatment As Usual. <i>The American Journal of Psychiatry</i> , 165(5), 631–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18347003">http://www.ncbi.nlm.nih.gov/pubmed/18347003</a>	<b>Borderline personality disorder</b>	<b>Mentalization based treatment (MBT;</b> Bateman & Fonagy, 2004)	<b>TAU</b>
17.	<b>Bateman, A., &amp; Fonagy, P. (2009).</b> Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1355-1364. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19833787">http://www.ncbi.nlm.nih.gov/pubmed/19833787</a>	<b>Borderline personality disorder</b>	<b>Mentalization based treatment (MBT;</b> Bateman & Fonagy, 2004)	<b>Structured Clinical Management (SCM)</b>
18.	<b>Ben-Itzhak, S., Bluvstein, I., Schreiber, S., Aharonov-Zaig, I., Maor, M., Lipnik, R., &amp; Bloch, M. (2012).</b> The Effectiveness of Brief Versus Intermediate Duration Psychodynamic Psychotherapy in the Treatment of Adjustment Disorder. <i>Journal of Contemporary Psychotherapy</i> , 42(4), 249–256.	<b>Adjustment disorder</b>	<b>Brief Psychodynamic Psychotherapy</b> (12 sessions; Malan, 1979)	<b>Brief Psychodynamic Psychotherapy</b> (1 yrs; Malan, 1979)

	<a href="http://www.springerlink.com/index/10.1007/s10879-012-9208-6">http://www.springerlink.com/index/10.1007/s10879-012-9208-6</a>			
19.	<b>Beresnevaite, M.</b> (2000) Exploring the benefits of group psychotherapy in reducing alexithymia in coronary heart disease patients: a preliminary study. <i>Psychotherapy and Psychosomatics</i> , 69, 117–122. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10773774?dopt=Abstract">http://www.ncbi.nlm.nih.gov/pubmed/10773774?dopt=Abstract</a>	<b>Alexithymia</b> (in patients with coronary heart disease)	<b>Group psychotherapy</b> (no reference)	<b>Psychoeducation</b>
20.	<b>Beutel, M. E., Dippel, A., Szczepanski, M., Thiede, R., &amp; Wiltink, J.</b> (2006). Mid-Term Effectiveness of Behavioral and Psychodynamic Inpatient Treatments of Severe Obesity Based on a Randomized Study. <i>Psychotherapy and Psychosomatics</i> , 75(6), 337–345. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17053334?dopt=Abstract">http://www.ncbi.nlm.nih.gov/pubmed/17053334?dopt=Abstract</a>	<b>Severe obesity</b>	<b>Individual in-patient PDT</b> (Local program, no reference)	<b>In-patient CBT</b> (Local program, no reference)
21.	<b>Beutel, M. E., Scheurich, V., Knebel, A., Michal, M., Wiltink, J., Graf-Morgenstern, M., ... Subic-Wrana, C.</b> (2013). Implementing panic-focused psychodynamic psychotherapy into clinical practice. <i>Canadian journal of psychiatry. Revue canadienne de psychiatrie</i> , 58(6), 326–34. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23768260">http://www.ncbi.nlm.nih.gov/pubmed/23768260</a>	<b>Panic disorder</b>	<b>Panic Focused Psychodynamic Psychotherapy</b> (PFPP; Milrod)	<b>CBT</b>
22.	<b>Beutel, M. E., Weissflog, G., Leuteritz, K., Wiltink, J., Haselbacher, a., Ruckes, C., ... Brahler, E.</b> (2013). Efficacy of short-term psychodynamic psychotherapy (STPP) with depressed breast cancer patients: results of a randomized controlled multicenter trial. <i>Annals of Oncology</i> , 1–7. <a href="http://annonc.oxfordjournals.org/cgi/doi/10.1093/annonc/mdt526">http://annonc.oxfordjournals.org/cgi/doi/10.1093/annonc/mdt526</a>  <b>Weißflog, G., Brähler, E., Leuteritz, K., Barthel, Y., Kuhnt, S., Wiltink, J., ... Beutel, M. E.</b> (2015). Does psychodynamic short-term psychotherapy for depressed breast cancer patients also improve fatigue? Results from a randomized controlled trial. <i>Breast Cancer Research and Treatment</i> . <a href="http://link.springer.com/10.1007/s10549-015-3494-0">http://link.springer.com/10.1007/s10549-015-3494-0</a>	<b>Depression</b> (in breast cancer patients)	<b>Supportive-expressive therapy</b> (Haselbacher, Barthel, Brähler et al., 2010; based on Luborsky, 1984)	<b>TAU</b>
23.	<b>Bloch, M., Meiboom, H., Lorberblatt, M., Bluvstein, I., Aharonov, I., &amp; Schreiber, S.</b> (2012). The effect of sertraline add-on to brief dynamic psychotherapy for the treatment of postpartum depression: a randomized, double-blind, placebo-controlled study. <i>The Journal of clinical psychiatry</i> , 73(2), 235–41. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22401479">http://www.ncbi.nlm.nih.gov/pubmed/22401479</a>	<b>Depression</b> (post-partum)	<b>Brief Dynamic Psychotherapy</b> (BDT; Malan, 1979) + <b>placebo meds</b>	<b>Brief Dynamic Psychotherapy</b> (BDT; Malan, 1979) + <b>Sertraline</b>
24.	<b>Bögels, S., Wijts, P., Oort, F., &amp; Sallaerts, S. J.</b> (2014). Psychodynamic psychotherapy versus cognitive-behavior therapy for social anxiety disorder: An efficacy and partial effectiveness trial. <i>Depression and Anxiety</i> , 31(5), 363–373. <a href="http://doi.wiley.com/10.1002/da.22246">http://doi.wiley.com/10.1002/da.22246</a>	<b>Social anxiety disorder (SAD)</b>	<b>STDP</b> (Malan, 1979; also Luborsky 1984)	<b>CBT</b> (Clark & Wells, 1995)
25.	<b>Bressi, C., Porcellana, M., Marinaccio, P. M., Nocito, E. P., &amp; Magri, L.</b> (2010). Short-term psychodynamic psychotherapy versus	<b>Mixed sample</b> (depression and anxiety)	<b>Brief Dynamic Psychotherapy</b> (BDT; Malan, 1976; Osimo &	<b>TAU</b>

	treatment as usual for depressive and anxiety disorders: a randomized clinical trial of efficacy. <i>The Journal of Nervous and Mental Disease</i> , 198(9), 647–52. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20823726">http://www.ncbi.nlm.nih.gov/pubmed/20823726</a>		Malan, 1992)	
26.	<b>Brockman</b> , B., Poynton, A., Ryle, A., & Watson, J. P. (1987). Effectiveness of time-limited therapy carried out by trainees. Comparison of two methods. <i>Br J Psychiatry</i> , 151, 602-610. <a href="http://bjp.rcpsych.org/content/151/5/602">http://bjp.rcpsych.org/content/151/5/602</a>	<b>Mixed sample</b>	<b>Interpretive therapy</b> (Mann & Goldman, 1982)	<b>Cognitive-analytic therapy</b> (CAT; Ryle, 1979; 1982)
27.	<b>Brodsky</b> , H., & Andrews, G. (1983). Brief psychotherapy in family practice. A controlled prospective intervention trial. <i>Br J Psychiatry</i> , 143, 11-19. <a href="http://bjp.rcpsych.org/content/143/1/11.short">http://bjp.rcpsych.org/content/143/1/11.short</a>	<b>Mixed sample</b>	<b>STDP</b> (Balant, Malan & Sifneos)	<b>1. Family practitioner</b> <b>2. No treatment</b>
28.	<b>Brom</b> , D., Kleber, R. J., & Defares, P. B. (1989). Brief psychotherapy for posttraumatic stress disorders. <i>J Consult Clin Psychol</i> , 57, 607-612. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2571625">http://www.ncbi.nlm.nih.gov/pubmed/2571625</a>	<b>PTSD</b>	<b>Brief Psychodynamic Therapy</b> (Horowitz, 1976)	<b>1. Trauma Desensitization</b> (no specific reference) <b>2. Hypnotherapy</b> (no specific reference)
29.	<b>Budman</b> , S. H., Demby, A., Redondo, J. P., Hannan, M., Feldstein, M., & Ring, J., et al. (1988). Comparative outcome in time-limited individual and group psychotherapy. <i>Int J Group Psychother</i> , 38, 63-86. <a href="http://www.tandfonline.com/doi/abs/10.1080/00207284.1988.11491085">http://www.tandfonline.com/doi/abs/10.1080/00207284.1988.11491085</a>	<b>Mixed sample</b>	<b>Time-limited individual psychotherapy</b> (Mann, 1973)	<b>Time-limited group psychotherapy</b> (Budman & Bennett, 1983)
30.	<b>Burnand</b> , Y., Andreoli, A., Kolatte, E., Venturini, A., & Rosset, N. (2002). Psychodynamic psychotherapy and clomipramine in the treatment of major depression. <i>Psychiatr Serv</i> , 53, 585-590. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11986508">http://www.ncbi.nlm.nih.gov/pubmed/11986508</a>	<b>Depression</b>	<b>PDT + clomipramine</b> , (unclear which model, Safran & Muran is mentioned among references)	<b>Supportive care + clomipramine</b>
31.	<b>Burns</b> , A., Guthrie, E., Marino-Francis, F., Busby, C., Morris, J., Russell, E., ... Byrne, J. (2005). Brief psychotherapy in Alzheimer's disease. <i>British Journal Of Psychiatry</i> , 187(2), 143–147. <a href="http://bjp.rcpsych.org/content/187/2/143.short">http://bjp.rcpsych.org/content/187/2/143.short</a>	<b>Alzheimers disease</b>	<b>Psychodynamic-Interpersonal therapy</b> (Hobson, 1985)	<b>TAU</b>
32.	<b>Chavooshi</b> , B., Mohammadkhani, P., & Dolatshahee, B. (2016). A Randomized Double-Blind Controlled Trial Comparing Davanloo's Intensive Short-Term Dynamic Psychotherapy as Internet-Delivered Versus Treatment as Usual for Medically Unexplained Pain: A Six-Month Pilot Study. <i>Psychosomatics</i> , 57(3), 292–300. <a href="http://linkinghub.elsevier.com/retrieve/pii/S033318216000025">http://linkinghub.elsevier.com/retrieve/pii/S033318216000025</a>	<b>Medically unexplained pain (MUP)</b>	<b>Internet-delivered ISTDP</b> (Abbass, 2015)	<b>TAU</b>
33.	<b>Chavooshi</b> , B., Mohammadkhani, P., & Dolatshahee, B. (2016). Efficacy of Intensive Short-Term Dynamic Psychotherapy for Medically Unexplained Pain: A Pilot Three-Armed Randomized Controlled Trial Comparison with Mindfulness-Based Stress Reduction. <i>Psychotherapy and Psychosomatics</i> , 123–125. <a href="http://www.karger.com/?doi=10.1159/000441698">http://www.karger.com/?doi=10.1159/000441698</a>	<b>Medically unexplained pain (MUP)</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Abbass, 2015)	<b>1. Mindfulness-based stress reduction (MBSR; no reference)</b> <b>2. TAU</b>

34.	<p><b>Chavooshi, B., Mohammadkhani, P., &amp; Dolatshahee, B. (2016).</b> Telemedicine vs. in-person delivery of intensive short-term dynamic psychotherapy for patients with medically unexplained pain: A 12-month randomized, controlled trial. <i>Journal of Telemedicine and Telecare, 0</i>(0), 1–9. <a href="http://jtt.sagepub.com/lookup/doi/10.1177/1357633X15627382">http://jtt.sagepub.com/lookup/doi/10.1177/1357633X15627382</a></p>	<b>Medically unexplained pain (MUP)</b>	<b>Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)</b>	<b>Internet-delivered ISTDP (ID-ISTDP)</b>
35.	<p><b>Clarici, A., Pellizzoni, S., Guaschino, S., Alberico, S., Bembich, S., Giuliani, R., ... Panksepp, J. (2015).</b> Intranasal administration of oxytocin in postnatal depression: implications for psychodynamic psychotherapy from a randomized double-blind pilot study. <i>Frontiers in Psychology, 06</i>, 1–10. <a href="http://journal.frontiersin.org/article/10.3389/fpsyg.2015.00426/abstract">http://journal.frontiersin.org/article/10.3389/fpsyg.2015.00426/abstract</a></p>	<b>Depression (post-partum)</b>	<b>Psychodynamic approach (non-manualized?) + oxytocine</b>	<b>Psychodynamic approach (non-manualized?) + placebo</b>
36.	<p><b>Clarkin, J. F., Levy, K. N., Lenzenweger, M. F., &amp; Kernberg, O. F. (2007).</b> Evaluating three treatments for borderline personality disorder: a multiwave study. <i>Am J Psychiatry, 164</i>, 922-928. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17541052">http://www.ncbi.nlm.nih.gov/pubmed/17541052</a></p> <p>Levy, K. N., Meehan, K. B., Kelly, K. M., Reynoso, J. S., Weber, M., Clarkin, J. F., &amp; Kernberg, O. F. (2006). Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. <i>Journal of Consulting and Clinical Psychology, 74</i>(6), 1027–40. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17154733">http://www.ncbi.nlm.nih.gov/pubmed/17154733</a></p>	<b>Borderline personality disorder</b>	<b>Transference-Focused Psychotherapy (TFP; Kernberg)</b>	<p><b>1. DBT (Linehan, 1991)</b></p> <p><b>2. Supportive treatment (Rockland, 1992)</b></p>
37.	<p><b>Connolly Gibbons, M. B., Thompson, S. M., Scott, K., Schauble, L. a, Mooney, T., Thompson, D., ... Crits-Christoph, P. (2012).</b> Supportive-expressive dynamic psychotherapy in the community mental health system: A pilot effectiveness trial for the treatment of depression. <i>Psychotherapy (Chicago, Ill.), 49</i>(3), 303–16. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22962971">http://www.ncbi.nlm.nih.gov/pubmed/22962971</a></p>	<b>Depression</b>	<b>Supportive-expressive therapy (manual adapted from Luborsky, 1984)</b>	<b>TAU</b>
38.	<p><b>Connolly Gibbons, M. B., Gallop, R., Thompson, D., Luther, D., Crits-Christoph, K., Jacobs, J., ... Crits-Christoph, P. (2016).</b> Comparative effectiveness of Cognitive Therapy and Dynamic Psychotherapy for Major Depressive Disorder in a community mental health setting. <i>JAMA Psychiatry, 173</i>(1), 1–8. <a href="http://doi.org/10.1001/jamapsychiatry.2016.1720">http://doi.org/10.1001/jamapsychiatry.2016.1720</a></p>	<b>Depression</b>	<b>Supportive-expressive therapy (Book, 1998; Luborsky, 1984)</b>	<b>CBT (Beck, 1995)</b>
39.	<p><b>Cooper, P. J., Murray, L., Wilson, A., &amp; Romaniuk, H. (2003).</b> Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. <i>Br J Psychiatry, 182</i>, 412-419. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12724244">http://www.ncbi.nlm.nih.gov/pubmed/12724244</a></p>	<b>Depression (post-partum)</b>	<b>Psychodynamic therapy (Cramer et al, 1990; Stern, 1995)</b>	<p><b>1. TAU</b></p> <p><b>2. Non-directive counseling (Holden et al, 1989)</b></p> <p><b>3. Cognitive-Behavioural Therapy (Hawton et al,</b></p>

	Murray, L., Cooper, P. J., Wilson, A., & Romaniuk, H. (2003). Controlled trial of the short- and long-term effect of psychological treatment of post- partum depression: 2. Impact on the mother-child relationship and child outcome. <i>British Journal of Psychiatry</i> , 182, 420–7. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12724245">http://www.ncbi.nlm.nih.gov/pubmed/12724245</a>			1989)
40.	<b>Covi, L., &amp; Lipman, R. S.</b> (1987). Cognitive behavioral group psychotherapy combined with imipramine in major depression. <i>Psychopharmacol Bull</i> , 23, 173-176. <a href="http://www.ncbi.nlm.nih.gov/pubmed/3602315">http://www.ncbi.nlm.nih.gov/pubmed/3602315</a>	<b>Depression</b>	<b>Traditional Group Therapy</b> (based on “psychodynamic-Interpersonal” theories but with no reference)	<b>1. CBT group therapy</b> (Beck, 1979; Covi, 1982) <b>2. CBT group therapy + Imipramine</b>
41.	<b>Covi, L., Lipman, R. S., Derogatis, L. R., Smith, J. E., &amp; Pattison, J. H.</b> (1974). Drugs and group psychotherapy in neurotic depression. <i>Am J Psychiatry</i> , 131, 191-198. <a href="http://ajp.psychiatryonline.org/doi/abs/10.1176/ajp.131.2.191">http://ajp.psychiatryonline.org/doi/abs/10.1176/ajp.131.2.191</a>	<b>Depression (chronic)</b>	<b>Psychodynamically oriented group therapy</b> (no reference) + <b>medication or placebo</b>	<b>Brief supportive contact + medication or placebo</b>
42.	<b>Creed, F., Fernandes, L., Guthrie, E., Palmer, S., Ratcliffe, J., &amp; Read, N., et al.</b> (2003). The cost-effectiveness of psychotherapy and paroxetine for severe irritable bowel syndrome. <i>Gastroenterology</i> , 124, 303-317. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12557136">http://www.ncbi.nlm.nih.gov/pubmed/12557136</a>  <b>Hyphantis, T., Guthrie, E., Tomenson, B., &amp; Creed, F.</b> (2009). Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome. <i>Pain</i> , 145(1-2), 196–203. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19643544">http://www.ncbi.nlm.nih.gov/pubmed/19643544</a>	<b>Irritable bowel syndrome</b>	<b>Psychodynamic-Interpersonal Therapy</b> (PIT; Hobson, 1985)	<b>TAU/SSRI</b>
43.	<b>Crits-Christoph, P., Gibbons, M. B. C., Narducci, J., Schamberger, M., &amp; Gallop, R.</b> (2005). Interpersonal Problems and the Outcome of Interpersonally Oriented Psychodynamic Treatment of GAD. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 42(2), 211–224. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.2.211">http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.2.211</a>	<b>Generalized anxiety disorder (GAD)</b>	<b>Supportive Expressive Therapy</b> (Luborsky (1984)	<b>Supportive therapy</b> (Borkovec & Matthews, 1988; Borkovec & Costello, 1993)
44.	<b>Crits-Christoph, P., Siqueland, L., Blaine, J., Frank, A., Luborsky, L., &amp; Onken, L. S., et al.</b> (1999). Psychosocial treatments for cocaine dependence: National Institute on Drug Abuse Collaborative Cocaine Treatment Study. <i>Arch Gen Psychiatry</i> , 56, 493-502. <a href="http://archpsyc.jamanetwork.com/article.aspx?articleid=1673777">http://archpsyc.jamanetwork.com/article.aspx?articleid=1673777</a>  <b>Crits-Christoph, P., Gibbons, M. B. C., Gallop, R., Ring-Kurtz, S., Barber, J. P., Worley, M., ... Hearon, B.</b> (2008). Supportive-Expressive Psychodynamic Therapy for Cocaine Dependence: A Closer Look. <i>Psychoanalytic Psychology : The Official Journal of the Division of Psychoanalysis, American Psychological Association, Division 39</i> , 25(3), 483–498. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2786223&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2786223&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Cocaine dependence</b>	<b>Supportive Expressive Therapy</b> (Luborsky (1984) + <b>group drug counseling</b> (Mercer & Woody, 1992)	<b>1. Cognitive therapy</b> (Beck, 1993) + <b>group drug counseling</b> (Mercer & Woody, 1992)  <b>2. Group drug counseling alone</b>

45.	<p>Cummings, J. G., &amp; Wittenberg, J. V. (2008). Supportive expressive therapy - Parent child version: an exploratory study. <i>Psychotherapy Theory, Research, Practice, Training</i>, 45, 148-164.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/22122414">http://www.ncbi.nlm.nih.gov/pubmed/22122414</a></p>	<p><b>Oppositional defiant disorder/disruptive behavior disorder</b> (children)</p>	<p><b>Supportive Expressive Therapy – Parent Child</b> (SET-PC; Wittenberg, adapted from Luborsky, 19854)</p>	<p><b>The Incredible Years Parenting Program</b></p>
46.	<p>Dare, C., Eisler, I., Russell, G., Treasure, J., &amp; Dodge, L. (2001). Psychological therapies for adults with anorexia nervosa: randomised controlled trial of out-patient treatments. <i>Br J Psychiatry</i>, 178, 216-221.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/11230031">http://www.ncbi.nlm.nih.gov/pubmed/11230031</a></p>	<p><b>Anorexia</b></p>	<p><b>Focal psychoanalytic psychotherapy</b> (Dare, 1995; Malan, 1979)</p>	<p><b>Cognitive Analytic Therapy</b> (Ryle, 1990; Treasure et al, 1995)</p> <p><b>Family therapy</b> (no specific reference)</p>
47.	<p>de Jonghe, F., Hendriksen, M., van Aalst, G., Kool, S., Peen, J., &amp; Van, R., et al. (2004). Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. <i>British Journal of Psychiatry</i>, 185, 37-45.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/15231554">http://www.ncbi.nlm.nih.gov/pubmed/15231554</a></p>	<p><b>Depression</b></p>	<p><b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, 2005)</p>	<p><b>Short Psychodynamic Supportive Psychotherapy + antidepressants</b></p>
48.	<p>de Jonghe, F., Kool, S., van Aalst, G., Dekker, J., &amp; Peen, J. (2001). Combining psychotherapy and antidepressants in the treatment of depression. <i>Journal of Affective Disorders</i>, 64(2-3), 217–29.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/11313088">http://www.ncbi.nlm.nih.gov/pubmed/11313088</a></p>	<p><b>Depression</b></p>	<p><b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, 2005) + <b>medication</b></p>	<p><b>Medication alone</b></p>
49.	<p>Dekker, J. J. M., Koelen, J. A., Van, H. L., Schoevers, R. A., Peen, J., Hendriksen, M., ... De Jonghe, F. (2008). Speed of action: the relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. <i>Journal of Affective Disorders</i>, 109(1-2), 183–188.  <a href="http://linkinghub.elsevier.com/retrieve/pii/S0165032707003588">http://linkinghub.elsevier.com/retrieve/pii/S0165032707003588</a></p> <p>Dekker, J., Van, H. L., Hendriksen, M., Koelen, J., Schoevers, R. A., Kool, S., ... Peen, J. (2013). What is the best sequential treatment strategy in the treatment of depression? Adding pharmacotherapy to psychotherapy or vice versa? <i>Psychotherapy and Psychosomatics</i>, 82(2), 89–98.  <a href="http://www.karger.com/doi/10.1159/000341177">http://www.karger.com/doi/10.1159/000341177</a></p>	<p><b>Depression</b></p>	<p><b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, 2005)</p>	<p><b>Pharmacotherapy</b></p>
50.	<p>Dekker, J., Molenaar, P. J., Kool, S., Van Aalst, G., Peen, J., &amp; de Jonghe, F. (2005). Dose–effect relations in time-limited combined psycho-pharmacological treatment for depression. <i>Psychological Medicine</i>, 35(1), 47–58.  <a href="http://www.journals.cambridge.org/abstract/S0033291704002685">http://www.journals.cambridge.org/abstract/S0033291704002685</a></p>	<p><b>Depression</b></p>	<p><b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, 2005) <b>8 sessions + medication</b></p>	<p><b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, 2005) <b>16 sessions + medication</b></p>
51.	<p>Doering, S., Hörz, S., Rentrop, M., Fischer-Kern, M., Schuster, P., &amp; Benecke, C., et al. (2010). Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: randomised controlled trial. <i>Br J Psychiatry</i>, 196, 389-395.</p>	<p><b>Borderline personality disorder</b></p>	<p><b>Transference-focused psychotherapy</b> (TFP; Kernberg)</p>	<p><b>TAU</b> (experienced therapists)</p>



	<a href="http://www.ncbi.nlm.nih.gov/pubmed/20435966">http://www.ncbi.nlm.nih.gov/pubmed/20435966</a>			
52.	<p><b>Driessen, E., Van, L. H., Don, F. J., Peen, J., Kool, S., Westra, D., ... Dekker, J. J. M. (2013).</b> The Efficacy of Cognitive-Behavioral Therapy and Psychodynamic Therapy in the Outpatient Treatment of Major Depression: A Randomized Clinical Trial. <i>American Journal of Psychiatry</i>, 170(9), 1041-1050 <a href="http://ajp.psychiatryonline.org/doi/abs/10.1176/appi.ajp.2013.12070899">http://ajp.psychiatryonline.org/doi/abs/10.1176/appi.ajp.2013.12070899</a></p> <p>Driessen, E., Van, H. L., Peen, J., Don, F. J., Kool, S., Westra, D., ... Dekker, J. J. (2014). Therapist-rated outcomes in a randomized clinical trial comparing cognitive behavioral therapy and psychodynamic therapy for major depression. <i>Journal of Affective Disorders</i>, 170C, 112–118. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25240140">http://www.ncbi.nlm.nih.gov/pubmed/25240140</a></p>	<b>Depression</b>	<b>Short-term psychodynamic supportive psychotherapy</b> (de Jonghe, 2005)	<b>Cognitive-Behavior Therapy (CBT;</b> Molenaar et al., 2009)
53.	<p><b>Durham, R. C., Murphy, T., Allan, T., Richard, K., Treliving, L. R., &amp; Fenton, G. W., et al. (1994).</b> Cognitive therapy, analytic psychotherapy and anxiety management training for generalised anxiety disorder. <i>Br J Psychiatry</i>, 165, 315-323. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7994500">http://www.ncbi.nlm.nih.gov/pubmed/7994500</a></p>	<b>Generalized anxiety disorder (GAD)</b>	<b>Analytic-based psychotherapy</b> (no specific reference)	<b>Cognitive Therapy / Anxiety Management training</b> (Beck & Emery, 1985)
54.	<p><b>Emmelkamp, P. M. G., Benner, A., Kuipers, A., Feiertag, G. A., Koster, H. C., &amp; Van Apeldoorn, F. J., et al. (2006).</b> Comparison of brief dynamic and cognitive-behavioural therapies in avoidant personality disorder. <i>Br J Psychiatry</i>, 189, 60-64. <a href="http://www.ncbi.nlm.nih.gov/pubmed/16816307">http://www.ncbi.nlm.nih.gov/pubmed/16816307</a></p>	<b>Avoidant personality disorder</b>	<b>Brief Dynamic Therapy (BDT;</b> Malan, 1979; also reference to Luborsky, 1984)	<b>CBT</b> (Beck & Freeman, 1990; Emmelkamp et al, 1992)
55.	<p><b>Fairburn, G., Kirk, J., Connor, M. O., &amp; Cooper, P. J. (1986).</b> A comparison of two psychological treatments for bulimia. <i>Behaviour research and therapy</i>, 24(6), 629–643. <a href="http://www.sciencedirect.com/science/article/pii/0005796786900586">http://www.sciencedirect.com/science/article/pii/0005796786900586</a></p>	<b>Bulimia</b>	<b>Short-term Focal Psychotherapy</b> (Rosen, 1979)	<b>CBT</b> (Fairburn, 1985)
56.	<p><b>Faramarzi, M., Azadfallah, P., Book, H. E., Tabatabaei, K. R., Taheri, H., &amp; Shokri-Shirvani, J. (2013).</b> A randomized controlled trial of brief psychoanalytic psychotherapy in patients with functional dyspepsia. <i>Asian journal of psychiatry</i>, 6(3), 228–34. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23642981">http://www.ncbi.nlm.nih.gov/pubmed/23642981</a></p> <p>Faramarzi, M., &amp; Book, H. E. (2015). The Effect of Psychotherapy in Improving Physical and Psychiatric Symptoms in Patients with Functional Dyspepsia, <i>Iran Journal of Psychiatry</i>, 43–49. <a href="http://www.ncbi.nlm.nih.gov/pubmed/26005480">http://www.ncbi.nlm.nih.gov/pubmed/26005480</a></p>	<b>Functional dyspepsia</b>	<b>Brief psychodynamic therapy</b> (Book, 1998) based on Luborsky (1984) + standard medical therapy	<b>TAU</b> (medical therapy)
57.	<p><b>Fonagy, P., Rost, F., Carlyle, J.-A., Mcpherson, S., Thomas, R., Fearon, R. M. P., ... Taylor, D. (2015).</b> Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression : the Tavistock</p>	<b>Depression</b> (treatment resistant)	<b>Long-Term Psychoanalytic PSYchotherapy (LTPP;</b> Taylor, 2015)	<b>TAU</b>

	Adult Depression Study (TADS ). <i>World Psychiatry</i> , 14(3), 312-321 <a href="http://www.ncbi.nlm.nih.gov/pubmed/26407787">http://www.ncbi.nlm.nih.gov/pubmed/26407787</a>			
58.	<b>Fonagy, P., Slead, M., &amp; Baradon, T.</b> (2016). Randomized Controlled Trial of Parent-Infant Psychotherapy for Parents With Mental Health Problems and Young Infants. <i>Infant Mental Health Journal</i> , 37(2), 97-114 <a href="http://doi.wiley.com/10.1002/imhj.21553">http://doi.wiley.com/10.1002/imhj.21553</a>	<b>Parents with mental health problems</b>	<b>Parent-infant psychotherapy (PIP)</b>	<b>Standard care</b>
59.	<b>Gallagher-Thompson, D., &amp; Steffen, A. M.</b> (1994). Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers. <i>J Consult Clin Psychol</i> , 62, 543-549. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8063980">http://www.ncbi.nlm.nih.gov/pubmed/8063980</a>	<b>Depression</b> (among family caregivers)	<b>Brief Psychodynamic Therapy</b> (Mann, 1973)	<b>CBT</b> (Beck et al., 1979; Lewinsohn et al, 1985; Gallagher et al 1981)
60.	<b>Gallagher, D. E., &amp; Thompson, L. W.</b> (1982). Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. <i>Psychotherapy: Theory, Research, and Practice</i> , 482-490. <a href="http://psycnet.apa.org/journals/pst/19/4/482">http://psycnet.apa.org/journals/pst/19/4/482</a>	<b>Depression</b>	<b>Brief relational/insight psychotherapy</b> (Bellak & Small, 1965)	<b>Behavioral therapy / Cognitive therapy</b> (Lewinsohn et al, 1976; Gallagher et al., 1981)
61.	<b>Garner, D. M., Rockert, W., Davis, R., Garner, M. V., Olmsted, M. P., &amp; Eagle, M., et al.</b> (1993). Comparison of cognitive-behavioral and supportive-expressive therapy for bulimia nervosa. <i>Am J Psychiatry</i> , 150, 37-46. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8417578">http://www.ncbi.nlm.nih.gov/pubmed/8417578</a>	<b>Bulimia</b>	<b>Supportive-expressive psychotherapy</b> (Luborsky, 1984)	<b>CBT</b> (Fairburn, 1985)
62.	<b>Gelder, M. G., Marks, I. M., &amp; Wolff, H. H.</b> (1967). Desensitization and psychotherapy in the treatment of phobic states: a controlled inquiry. <i>British Journal of Psychiatry</i> , 113(494), 53-81. <a href="http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.113.494.53">http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.113.494.53</a>	<b>Phobic states</b>	<b>Psychotherapy</b> (no reference)	<b>1. Desensitization</b> (Wolpe) <b>2. Group psychotherapy</b> (Foulkes)
63.	<b>Gersons, B. P., Carlier, I. V, Lamberts, R. D., &amp; van der Kolk, B. a.</b> (2000). Randomized clinical trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder. <i>Journal of Traumatic Stress</i> , 13(2), 333-47. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10838679">http://www.ncbi.nlm.nih.gov/pubmed/10838679</a>	<b>PTSD</b>	<b>Brief eclectic psychotherapy (BEP;</b> Gersons & Carlier, 1994; Manual includes both CBT and PDT components)	<b>Waitlist</b>
64.	<b>Ghorbani, N., Dadsetan, P., Ejei, J., &amp; Motiyan, H.</b> (2000). The consequences of overcoming resistance and emotional disclosure on lymphocyte T-helper and T-suppressor and psychological pathology. <i>Journal of Psychology (Persian)</i> , 3, 368-89.	<b>Students</b>	<b>ISTDP</b> (Davanloo, 1990)	<b>Simple verbal disclosure groups (SVD)</b>
65.	<b>Giesen-Bloo, J., van Dyck, R., Spinhoven, P., van Tilburg, W., Dirksen, C., &amp; van Asselt, T., et al.</b> (2006). Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. <i>Arch Gen Psychiatry</i> , 63, 649-658. <a href="http://www.ncbi.nlm.nih.gov/pubmed/16754838">http://www.ncbi.nlm.nih.gov/pubmed/16754838</a>  van Asselt, A. D. I., Dirksen, C. D., Arntz, A., Giesen-Bloo, J. H., van Dyck, R., Spinhoven, P.,	<b>Borderline personality disorder</b>	<b>Transference focused psychotherapy (TFP;</b> Clarkin, Yeomans, & Kernberg, 1999; Yeomans, Clarkin, & Kernberg, 2002)	<b>Schema-focused therapy</b> (Young, 1994; Young et al, 2003)

	<p>... Severens, J. L. (2008). Out-patient psychotherapy for borderline personality disorder: cost-effectiveness of schema-focused therapy v. transference-focused psychotherapy. <i>The British Journal of Psychiatry: The Journal of Mental Science</i>, 192(6), 450–7.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/18515897">http://www.ncbi.nlm.nih.gov/pubmed/18515897</a></p>			
66.	<p><b>Gilboa-Schechtman, E., Foa, E. B., Shafran, N., Aderka, I. M., Powers, M. B., Rachamim, L., ... Apter, A. (2010).</b> Prolonged exposure versus dynamic therapy for adolescent PTSD: a pilot randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i>, 49(10), 1034–42.  <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2945239&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2945239&amp;tool=pmcentrez&amp;rendertype=abstract</a></p>	<b>PTSD (adolescents)</b>	<b>Time-limited dynamic therapy (TLDP-A; Shefler et al, 1995)</b>	<b>Prolonged exposure (Foa)</b>
67.	<p><b>Gowers, S., Norton, K., Halek, C., &amp; Crisp, A. H. (1994).</b> Outcome of outpatient psychotherapy in a random allocation treatment study of anorexia nervosa. <i>Int J Eat Disord</i>, 15, 165-177.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/8173562">http://www.ncbi.nlm.nih.gov/pubmed/8173562</a></p>	<b>Anorexia (adolescents and young women)</b>	<b>Short term individual + family interviews (Koss &amp; Butcher, 1986)</b>	<b>Assessment only</b>
68.	<p><b>Gregory, R. J., Chlebowski, S., Kang, D., Remen, A. L., Soderberg, M. G., &amp; Stepkovitch, J., et al. (2008).</b> A controlled trial of psychodynamic psychotherapy for co-occurring borderline personality disorder and alcohol use disorder. <i>Psychotherapy: Theory, Research, Practice, Training</i>, 45, 28-41.  <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.45.1.28">http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.45.1.28</a></p>	<b>Borderline personality disorder + alcohol use disorder</b>	<b>Dynamic deconstructive therapy (Gregory &amp; Remen)</b>	<b>TAU</b>
69.	<p><b>Grothe, C., Bering, R., SpieB, R., Luneburg, E., &amp; Fischer, G. (2003).</b> Mehrdimensionale Psychodynamische Traumatherapie (MPTT) Forschungsergebnisse zur Standardversion. <i>Zeitschrift fur Psychotraumatologie, Psychotherapiewissenschaft und Psychologische Medizin</i> 1(2), 27–43.</p>	<b>PTSD</b>	<b>Mehrdimensionale Psychodynamische Traumatherapie (MPTT; Fischer, 2000)</b>	<p><b>1. EMDR (Shapiro, 1995; 1998)</b></p> <p><b>2. Waitlist control (?)</b></p>
70.	<p><b>Gunderson, J. G., Frank, A. F., Katz, H. M., Vannicelli, M. L., Frosch, J. P., &amp; Knapp, P. H., et al. (1984).</b> Effects of psychotherapy in schizophrenia: II. Comparative outcome of two forms of treatment. <i>Schizophr Bull</i>, 10, 564-598.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/6151246">http://www.ncbi.nlm.nih.gov/pubmed/6151246</a></p>	<b>Schizophrenia</b>	<b>Exploratory, insight oriented psychotherapy (Adapted from Fromm-Reichmann)</b>	<b>Reality-adaptive, supportive psychotherapy</b>
71.	<p><b>Guthrie, E., Creed, F., Dawson, D., &amp; Tomenson, B. (1991).</b> A controlled trial of psychological treatment for the irritable bowel syndrome. <i>Gastroenterology</i>, 100, 450-457.  <a href="http://www.gastrojournal.org/article/0016-5085(91)90215-7/pdf">http://www.gastrojournal.org/article/0016-5085(91)90215-7/pdf</a></p> <p>Guthrie, E., Creed, F., Dawson, D., &amp; Tomenson, B. (1993). A randomised controlled trial of psychotherapy in patients with refractory Irritable Bowel Syndrome. <i>British Journal of Psychiatry</i>, 163, 315–21.  <a href="http://bjp.rcpsych.org/content/163/3/315">http://bjp.rcpsych.org/content/163/3/315</a></p>	<b>Irritable bowel syndrome</b>	<b>Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)</b>	<b>Supportive listening</b>
72.	<p><b>Guthrie, E., Kapur, N., Mackway-Jones, K.,</b></p>	<b>Self-poisoning</b>	<b>Psychodynamic-</b>	<b>TAU</b>

	Chew-Graham, C., Moorey, J., & Mendel, E., et al. (2001). Randomised controlled trial of brief psychological intervention after deliberate self poisoning. <i>BMJ</i> , 323, 135-138. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=34723&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=34723&amp;tool=pmcentrez&amp;rendertype=abstract</a>		<b>Interpersonal Therapy</b> (PIT; Hobson, 1985)	
73.	<b>Guthrie, E.</b> , Moorey, J., Margison, F., Barker, H., Palmer, S., & McGrath, G., et al. (1999). Cost-effectiveness of brief psychodynamic-interpersonal therapy in high utilizers of psychiatric services. <i>Arch Gen Psychiatry</i> , 56, 519-526. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10359466">http://www.ncbi.nlm.nih.gov/pubmed/10359466</a>	<b>Mixed sample</b>	<b>Psychodynamic-Interpersonal Therapy</b> (PIT; Hobson, 1985)	<b>TAU</b>
74.	<b>Hall, A.</b> , & Crisp, A. H. (1987). Brief psychotherapy in the treatment of anorexia nervosa: Outcome at one year. <i>British Journal of Psychiatry</i> , 151, 185-191. <a href="http://www.ncbi.nlm.nih.gov/pubmed/369108">http://www.ncbi.nlm.nih.gov/pubmed/369108</a>	<b>Anorexia</b>	<b>Combined individual and family psychotherapy</b> (no reference)	<b>Dietary advice</b>
75.	<b>Hamilton, J.</b> , Guthrie, E., Creed, F., Thompson, D., Tomenson, B., & Bennett, R., et al. (2000). A randomized controlled trial of psychotherapy in patients with chronic functional dyspepsia. <i>Gastroenterology</i> , 119, 661-669. <a href="http://linkinghub.elsevier.com/retrieve/pii/S0016508500145949">http://linkinghub.elsevier.com/retrieve/pii/S0016508500145949</a>	<b>Chronic Functional Dyspepsia</b>	<b>Psychodynamic-Interpersonal Therapy</b> (PIT; Hobson, 1985)	<b>Supportive therapy</b> (Guthrie, unpub.)
76.	<b>Hardy, G. E.</b> , Barkham, M., Shapiro, D. a, Stiles, W. B., Rees, a, & Reynolds, S. (1995). Impact of Cluster C personality disorders on outcomes of contrasting brief psychotherapies for depression. <i>Journal of consulting and clinical psychology</i> , 63(6), 997-1004. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8543722">http://www.ncbi.nlm.nih.gov/pubmed/8543722</a>	<b>Depression + personality disorder cluster C</b>	<b>Psychodynamic-Interpersonal Therapy</b> (PIT; Hobson, 1985)	<b>CBT</b> (Firth & Shapiro, 1985)
77.	<b>Hellerstein, D. J.</b> , Rosenthal, R. N., Pinsker, H., Samstag, L. W., Muran, J. C., & Winston, A., et al. (1998). A randomized prospective study comparing supportive and dynamic therapies. Outcome and alliance. <i>J Psychother Pract Res</i> , 7, 261-271. <a href="http://psycnet.apa.org/psycinfo/1998-12001-001">http://psycnet.apa.org/psycinfo/1998-12001-001</a>	<b>Personality disorders</b> (cluster C)	<b>ISTDP</b> (Davanloo, 1980)	<b>Brief Adaptive Therapy</b> (Winston?)
78.	<b>Hersen, M.</b> , Himmelhoch, J. M., Thase, M. E., & Bellack, A. S. (1984). Effects of social skill training, amitriptyline, and psychotherapy in unipolar depressed women. <i>Behavior Therapy</i> , 15(1), 21-40. <a href="http://linkinghub.elsevier.com/retrieve/pii/S005789484800398">http://linkinghub.elsevier.com/retrieve/pii/S005789484800398</a>	<b>Depression</b>	<b>Time-limited dynamic therapy</b> (no reference) + placebo	<b>1. Social skill training + placebo</b>  <b>2. Amitriptyline</b>
79.	<b>Hoffart, A.</b> , & Martinsen, E. W. (1990). Exposure-based integrated vs. Pure psychodynamic treatment of agoraphobic inpatients. <i>Psychotherapy: Theory, Research, Practice, Training, Vol 27(2)</i> , 210-218. <a href="http://psycnet.apa.org/psycinfo/1990-29285-001">http://psycnet.apa.org/psycinfo/1990-29285-001</a>	<b>Agoraphobia</b>	<b>Integrative behavioural-psychodynamic treatment</b>	<b>Psychodynamic oriented group therapy</b>
80.	<b>Høglend, P.</b> , Amlø, S., Marble, A., Bøgwald, K. P., Sørbye, O., & Sjaastad, M. C., et al. (2006). Analysis of the patient-therapist relationship in dynamic psychotherapy: an experimental	<b>Mixed sample</b>	<b>Psychodynamic psychotherapy with transference interpretation</b> (Høglend,	<b>Psychodynamic psychotherapy without transference interpretation</b> (Høglend,

	<p>study of transference interpretations. <i>Am J Psychiatry</i>, 163, 1739-1746.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/17012684">http://www.ncbi.nlm.nih.gov/pubmed/17012684</a></p> <p>Høglend, P., Bøgwald, K.-P., Amlø, S., Marble, A., Ulberg, R., Sjaastad, M. C., ... Johansson, P. (2008). Transference interpretations in dynamic psychotherapy: do they really yield sustained effects? <i>The American Journal of Psychiatry</i>, 165(6), 763–71.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/18413707">http://www.ncbi.nlm.nih.gov/pubmed/18413707</a></p>		1995)	1995)
81.	<p><b>Huber, D., Henrich, G., Gastner, J., &amp; Klug, G.</b> (2012). Must All Have Prizes? The Munich Psychotherapy Study. In <i>Psychodynamic Psychotherapy Research</i>.  <a href="http://link.springer.com/chapter/10.1007%2F978-1-60761-792-1_4#page-1">http://link.springer.com/chapter/10.1007%2F978-1-60761-792-1_4#page-1</a></p> <p>Huber, D., Huber, D., Zimmermann, J., Henrich, G., &amp; Klug, G. (2012). Comparison of cognitive-behaviour therapy with psychoanalytic and psychodynamic therapy for depressed patients – A three-year follow-up study. <i>Zeitschrift für Psychosomatic</i>, (58), 299–316.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/22987495">http://www.ncbi.nlm.nih.gov/pubmed/22987495</a></p> <p>Huber, D., Henrich, G., Clarkin, J., &amp; Klug, G. (2013). Psychoanalytic versus psychodynamic therapy for depression: a three-year follow-up study. <i>Psychiatry</i>, 76(2), 132–49.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/23631544">http://www.ncbi.nlm.nih.gov/pubmed/23631544</a></p>	<b>Depression</b>	<b>Psychoanalysis</b> (non-manualized)	<p>1. <b>LTTP</b> (non-manualized)</p> <p>2. <b>CBT</b> (non-manualized; patients not randomly assigned to this group)</p>
82.	<p><b>Hubschmid, M., Aybek, S., D, M., Maccaferri, G. E., D, M., Chocron, O., ... D, M. (n.d.)</b>. Efficacy of brief interdisciplinary psychotherapeutic intervention for motor conversion disorder and nonepileptic attacks. <i>General Hospital Psychiatry</i>, 37(5), 448–455. doi:10.1016/j.genhosppsych.2015.05.007  <a href="http://dx.doi.org/10.1016/j.genhosppsych.2015.05.007">http://dx.doi.org/10.1016/j.genhosppsych.2015.05.007</a></p>	<b>Motor conversion disorder and nonepileptic seizures</b>	<b>Interdisciplinary psychotherapeutic intervention</b> (Psychodynamic-Interpersonal therapy based on Guthrie, 1999)	<b>Standard care</b>
83.	<p><b>Jantschek, G., Zeitz, M., Wirsching, M., Klor, H. U., Studt, H. H., Rasenack, J., ... Disease, T. G. S. G. on P. I. in C.</b> (1999). Effect of psychotherapy on the course of Crohn's disease. <i>Psychosomatic Intervention in CD</i>, 1289–1296.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/9930393">http://www.ncbi.nlm.nih.gov/pubmed/9930393</a></p> <p>Keller, W., Pritsch, M., Von Wietersheim, J., Scheib, P., Osborn, W., &amp; Balck, F., et al. (2004). Effect of psychotherapy and relaxation on the psychosocial and somatic course of Crohn's disease: main results of the German Prospective Multicenter Psychotherapy Treatment study on Crohn's Disease. <i>J Psychosom Res</i>, 56, 687-696.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/15193965">http://www.ncbi.nlm.nih.gov/pubmed/15193965</a></p>	<b>Crohn's disease</b>	<b>PDT+relaxation+drug treatment</b> (no manual was used but reference to Luborsky, 1984)	<b>Drug treatment</b>
84.	<p><b>Johansson, R., Björklund, M., Hornborg, C., Karlsson, S., Hesser, H., Ljótsson, B., ... Andersson, G.</b> (2013). Affect-focused</p>	<b>Depression and/or anxiety</b>	<b>IPDT</b> (Affect-focused; based on Frederick, 2009)	<b>Online therapist support and clinical monitoring</b>

	psychodynamic psychotherapy for depression and anxiety through the Internet: a randomized controlled trial. <i>PeerJ</i> , 1, e102. <a href="https://peerj.com/articles/102">https://peerj.com/articles/102</a>			
85.	<b>Johansson, R., Ekbladh, S., Hebert, A., Lindström, M., Möller, S., Petitt, E., ... Andersson, G. (2012).</b> Psychodynamic Guided Self-Help for Adult Depression through the Internet: A Randomised Controlled Trial. (A. Bruce, Ed.) <i>PLoS ONE</i> , 7(5), <a href="http://dx.plos.org/10.1371/journal.pone.0038021">http://dx.plos.org/10.1371/journal.pone.0038021</a>	<b>Depression</b>	<b>IPDT (SUBGAP; based on Silverberg, 2005)</b>	<b>Psychoeducation + support</b>
86.	<b>Jørgensen, C. R., Freund, C., Bøye, R., Jordet, H., Andersen, D., &amp; Kjølbye, M. (2012).</b> Outcome of mentalization-based and supportive psychotherapy in patients with borderline personality disorder: a randomized trial. <i>Acta psychiatrica Scandinavica</i> , 1–13. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22897123">http://www.ncbi.nlm.nih.gov/pubmed/22897123</a>	<b>Borderline personality disorder</b>	<b>Mentalization-Based Therapy (MBT; Bateman &amp; Fonagy, 2001)</b>	<b>Supportive group therapy</b>
87.	<b>Klein, D. F., Zitrin, C. M., Woerner, M. G., &amp; Ross, D. C. (1983).</b> Treatment of Phobias. II. Behavior therapy and supportive psychotherapy: are there any specific ingredients? <i>Archives of General Psychiatry</i> , 139–145. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6130751">http://www.ncbi.nlm.nih.gov/pubmed/6130751</a>	<b>Phobias</b>	<b>Supportive therapy (dynamically oriented; no reference)</b>	<b>Behavior therapy (no reference)</b>
88.	<b>Kleinman, P. H., Woody, G. E., Todd, T. C., Millman, R. B., Kang, S. Y., &amp; Kemp, J., et al. (1990).</b> Crack and cocaine abusers in outpatient psychotherapy. <i>NIDA Res Monogr</i> , 104, 24-35.	<b>Cocaine abuse</b>	<b>Supportive-Expressive Therapy (Luborsky, 1984)</b>	<b>Family therapy + para professionally led group therapy</b>
89.	<b>Knekt, P., &amp; Lindfors, O. (2004).</b> A randomized trial of the effect of four forms of psychotherapy on depressive and anxiety disorders. Helsinki: Kela. <a href="https://helda.helsinki.fi/handle/10250/3315">https://helda.helsinki.fi/handle/10250/3315</a>  <b>Knekt, P., Lindfors, O., Ske, L. S. Ä., &amp; Virtala, E. S. A. (2013).</b> Randomized trial on the effectiveness of long- and short-term psychotherapy on psychiatric symptoms and working ability during a 5-year follow-up. <i>Nordic journal of psychiatry</i> , 67(1), 59-68. <a href="http://www.tandfonline.com/doi/full/10.3109/08039488.2012.680910">http://www.tandfonline.com/doi/full/10.3109/08039488.2012.680910</a>  <b>Knekt, P., Virtala, E., Härkänen, T., Vaarama, M., Lehtonen, J., &amp; Lindfors, O. (2016).</b> The outcome of short- and long-term psychotherapy 10 years after start of treatment. <i>Psychological Medicine</i> , 1–14. <a href="http://www.journals.cambridge.org/abstract/S0033291715002718">http://www.journals.cambridge.org/abstract/S0033291715002718</a>  <b>Lindfors, O., Knekt, P., Virtala, E., &amp; Laaksonen, M. a. (2012).</b> The effectiveness of solution-focused therapy and short- and long-term psychodynamic psychotherapy on self-concept during a 3-year follow-up. <i>The Journal of Nervous and Mental Disease</i> , 200(11), 946–53. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23124178">http://www.ncbi.nlm.nih.gov/pubmed/23124178</a>	<b>Depression and/or anxiety</b>	<b>STPP (Malan, 1976; Sifneos, 1978)</b>	<b>1. Solution-focused therapy (de Shazer et al, 1986)</b>  <b>2. LTPP (Gabbard, 2004)</b>

	<p>Lindfors, O., Knekt, P., Heinonen, E., Härkänen, T., Virtala, E., &amp; The Helsinki Psychotherapy Study Group. (2015). The effectiveness of short- and long-term psychotherapy on personality functioning during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 173, 31–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25462393">http://www.ncbi.nlm.nih.gov/pubmed/25462393</a></p> <p>Maljanen, T., Knekt, P., Lindfors, O., Virtala, E., Tillman, P., &amp; Härkänen, T. (2016). The cost-effectiveness of short-term and long-term psychotherapy in the treatment of depressive and anxiety disorders during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 190(July), 254–263. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22611089">http://www.ncbi.nlm.nih.gov/pubmed/22611089</a></p>			
90.	<p><b>Knijnik, D. Z., Blanco, C., Salum, G. A., Moraes, C. U., Mombach, C., Almeida, E., ... Eizirik, C. L.</b> (2008). A pilot study of clonazepam versus psychodynamic group therapy plus clonazepam in the treatment of generalized social anxiety disorder. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 23(8), 567–74. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18774274">http://www.ncbi.nlm.nih.gov/pubmed/18774274</a></p>	<b>Social anxiety disorder (SAD)</b>	<b>Psychodynamic group therapy (Malan, 1976) + clonazepam</b>	<b>Clonazepam</b>
91.	<p><b>Knijnik, D. Z., Kapczinski, F., Chachamovich, E., Margis, R., &amp; Eizirik, C. L.</b> (2004). Psychodynamic group treatment for generalized social phobia. <i>Revista Brasileira de Psiquiatria (São Paulo, Brazil : 1999)</i>, 26(2), 77–81. <a href="http://www.ncbi.nlm.nih.gov/pubmed/15517057">http://www.ncbi.nlm.nih.gov/pubmed/15517057</a></p>	<b>Social anxiety disorder (SAD)</b>	<b>Psychodynamic group therapy (Malan, 1976)</b>	<b>Credible placebo group</b>
92.	<p><b>Kompolti, K., Wilson, B., Stebbins, G., Bernard, B., &amp; Hinson, V.</b> (2013). Immediate vs. delayed treatment of psychogenic movement disorders with short-term psychodynamic psychotherapy: Randomized clinical trial. <i>Parkinsonism &amp; related disorders</i>, 7–10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24120952">http://www.ncbi.nlm.nih.gov/pubmed/24120952</a></p>	<b>Psychogenic movement disorders</b>	<b>ISTDP (Davanloo, 1980)</b>	<b>Neurological observation and support</b>
93.	<p><b>Kool, S., Dekker, J., &amp; Duijsens, I.</b> (2003). Efficacy of combined therapy and pharmacotherapy for depressed patients with or without personality disorders. <i>Harvard review of psychiatry</i>, 133–142. <a href="http://informahealthcare.com/doi/abs/10.1080/10673220303950">http://informahealthcare.com/doi/abs/10.1080/10673220303950</a></p>	<b>Depression</b>	<b>STDP (de Jonghe, 2005) + pharmacotherapy</b>	<b>Pharmacotherapy</b>
94.	<p><b>Kornblith, S., Rehm, L., W, O. M., &amp; Lamparski, D. M.</b> (1983). The contribution of self-reinforcement training and behavioral assignments to the efficacy of self-control therapy for depression. <i>Cogn Ther Res</i>, 7, 499–528. <a href="http://link.springer.com/10.1007/BF01172888">http://link.springer.com/10.1007/BF01172888</a></p>	<b>Depression</b>	<b>Psychodynamic group therapy (Wolf, 1965; Bernard &amp; Klein, 1977)</b>	<b>Self-reinforcement training and behavioral assignments (Rehm, 1977)</b>
95.	<p><b>Kramer, U., de Roten, Y., Perry, J. C., &amp; Despland, J.-N.</b> (2013). Change in defense mechanisms and coping patterns during the course of 2-year-long psychotherapy and psychoanalysis for recurrent depression: a pilot study of a randomized controlled trial.</p>	<b>Depression (recurrent)</b>	<p><b>Supportive-Expressive Therapy (Luborsky, 1984)</b></p> <p>All patients were also given pharmacological treatment</p>	<p><b>1. Cognitive therapy (Beck, 1995)</b></p> <p><b>2. Psychoanalysis (Robertson, 2002).</b></p>

	<i>The Journal of nervous and mental disease</i> , 201(7), 614–20. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23817160">http://www.ncbi.nlm.nih.gov/pubmed/23817160</a>			<b>3. Clinical management</b> (Csank, 2002; Novalis et al., 1993).
96.	<b>Lanza</b> , M. L., Anderson, J., Boisvert, C. M., LeBlanc, A., Fardy, M., & Steel, B., et al. (2002). Assaultive behavior intervention in the Veterans Administration: psychodynamic group psychotherapy compared to cognitive behavior therapy. <i>Perspect Psychiatr Care</i> , 38, 89-97. <a href="http://onlinelibrary.wiley.com/doi/10.1111/j.1744-6163.2002.tb00662.x/abstract">http://onlinelibrary.wiley.com/doi/10.1111/j.1744-6163.2002.tb00662.x/abstract</a>	<b>Male veterans with history of assault</b>	<b>Psychodynamic group psychotherapy</b> (no specific reference)	<b>Cognitive behavior group therapy</b> (No specific reference)
97.	<b>LaPonite</b> , K., & Rimm, D. C. (1980). Cognitive, assertive, and insight-oriented group therapies in the treatment of reactive depression in women. <i>Psychotherapy: Theory, Research &amp; Practice</i> , 17(3), 312–321. <a href="http://psycnet.apa.org/journals/pst/17/3/312">http://psycnet.apa.org/journals/pst/17/3/312</a>	<b>Depression</b>	<b>Insight-oriented group therapy</b> (No reference)	<b>1. Cognitive group therapy</b> (Beck, 19767) <b>2. Assertiveness training</b> (no reference)
98.	<b>Lau</b> , M., & Kristensen, E. (2007). Outcome of systemic and analytic group psychotherapy for adult women with history of intrafamilial childhood sexual abuse: a randomized controlled study. <i>Acta Psychiatr Scand</i> , 116, 96-104. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17650270">http://www.ncbi.nlm.nih.gov/pubmed/17650270</a>	<b>Women with childhood sexual abuse</b>	<b>Analytic group therapy</b> (Foulkes, 1986)	<b>Systemic group therapy</b> (Holme, 1999)
99.	<b>Lawson</b> , D. M. (2010). Comparing cognitive behavioral therapy and integrated cognitive behavioral therapy/psychodynamic therapy in group treatment for partner violent men. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 47(1), 122–133. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0018841">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0018841</a>	<b>Violent men</b>	<b>Integrative CBT- psychodynamic group therapy</b> (Strupp & Binder, 1984)	<b>CBT</b> (Lawson, 2001)
100.	<b>Leichsenring</b> , F., Masuhr, O., Jaeger, U., Rabung, S., Dally, A., Dumpelmann, M., ... Streeck, U. (2016). Psychoanalytic-Interactional Therapy versus Psychodynamic Therapy by Experts for Personality Disorders: A Randomized Controlled Efficacy-Effectiveness Study in Cluster B Personality Disorders. <i>Psychotherapy and Psychosomatics</i> , 71–80. <a href="http://www.karger.com/?doi=10.1159/000441731">http://www.karger.com/?doi=10.1159/000441731</a>	<b>Personality disorders (cluster B)</b>	<b>Psychoanalytic-Interactional Therapy</b> (Streeck, 2016)	<b>Psychodynamic therapy by experts</b>
101.	<b>Leichsenring</b> , F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E. (2013). Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder: A Multicenter Randomized Controlled Trial. <i>American Journal of Psychiatry</i> , 170(7), 759-767. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23680854">http://www.ncbi.nlm.nih.gov/pubmed/23680854</a>  <b>Leichsenring</b> , F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E. (2014). Long-Term Outcome of Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder. <i>American Journal of Psychiatry</i> , 171(10), 1074-1082.	<b>Social anxiety disorder (SAD)</b>	<b>Supportive-Expressive Therapy</b> (Luborsky, 1984)	<b>CBT</b> (Clark & Wells, 1995)



	<a href="http://www.ncbi.nlm.nih.gov/pubmed/25016974">http://www.ncbi.nlm.nih.gov/pubmed/25016974</a>			
102.	<p><b>Leichsenring, F., Salzer, S., Jaeger, U., Kächele, H., Kreische, R., &amp; Leweke, F., et al. (2009).</b> Short-Term Psychodynamic Psychotherapy and Cognitive-Behavioral Therapy in Generalized Anxiety Disorder: A Randomized, Controlled Trial. <i>Am J Psychiatry</i>, <i>166</i>, 875-881. <a href="http://ajp.psychiatryonline.org/article.aspx?articleID=101023&amp;atab=7">http://ajp.psychiatryonline.org/article.aspx?articleID=101023&amp;atab=7</a></p> <p>Salzer, S., Winkelbach, C., Leweke, F., Leibing, E., &amp; Leichsenring, F. (2011). Generalized Anxiety Disorder: 12-Month Follow-Up. <i>Canadian Journal of Psychiatry</i>, <i>56</i>(8), 1-10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24229481">http://www.ncbi.nlm.nih.gov/pubmed/24229481</a></p>	<b>Generalized anxiety disorder (GAD)</b>	<b>Supportive-Expressive Therapy</b> (Luborsky, 1984)	<b>CBT</b> (Borcovec & Rucio, 2001, Brown, 1993)
103.	<p><b>Lemma, A., &amp; Fonagy, P. (2013).</b> Feasibility study of a psychodynamic online group intervention for depression. <i>Psychoanalytic Psychology</i>, <i>30</i>(3), 367-380. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0033239">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0033239</a></p>	<b>Depression</b>	<b>Online Group Dynamic Interpersonal Therapy</b> (OLDIT; Fonagy & Target, 2008)	<b>Internet discussion group</b>
104.	<p><b>Lerner, A., Sigal, M., Bacalu, A., &amp; Gelkopf, M. (1992).</b> Short term versus long term psychotherapy in opioid dependence: A pilot study. <i>Israeli Journal of Psychiatry and Related Sciences</i>, <i>29</i>(2), 114-9. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1526756">http://www.ncbi.nlm.nih.gov/pubmed/1526756</a></p>	<b>Opiate dependence</b>	<b>Short-Term Dynamic Psychotherapy</b> (Luborsky, 1989; Mann, 1973; Davanloo, 1978)	<b>Long-Term Dynamic Psychotherapy</b> (Karasu, 1989)
105.	<p><b>Levy Berg, A., Sandell, R., &amp; Sandahl, C. (2009).</b> Affect-focused body psychotherapy in patients with generalized anxiety disorder: Evaluation of an integrative method. <i>Journal of Psychotherapy Integration</i>, <i>19</i>(1), 67-85. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0015324">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0015324</a></p>	<b>Generalized anxiety disorder (GAD)</b>	<b>Affect-Focused Body Psychotherapy</b> (Monsen & Monsen, 2000)	<b>TAU</b>
106.	<p><b>Liberman, R. P., &amp; Eckman, T. (1981).</b> Behavior therapy vs insight-oriented therapy for repeated suicide attempters. <i>Archives of General Psychiatry</i>, <i>38</i>(10), 1126-1130. <a href="http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1981.01780350060007">http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1981.01780350060007</a></p>	<b>Repeated suicide attempts</b>	<b>Insight-Oriented Psychotherapy</b>	<b>Behavior Therapy</b>
107.	<p><b>Lindauer, R. J. L., Gersons, B. P. R., Van Meijel, E. P. M., Blom, K., Carlier, I. V. E., Vrijlandt, I., &amp; Olff, M. (2005).</b> Effects of brief eclectic psychotherapy in patients with posttraumatic stress disorder: Randomized clinical trial. <i>Journal of Traumatic Stress</i>, <i>18</i>(3), 205-212. <a href="http://doi.wiley.com/10.1002/pts.20029">http://doi.wiley.com/10.1002/pts.20029</a></p>	<b>PTSD</b>	<b>Brief eclectic psychotherapy (BEP;</b> Gersons et al., 2004; Manual includes both CBT and PDT components)	<b>Waitlist</b>
108.	<p><b>Linnet, J., &amp; Jemec, G. B. (2001).</b> Anxiety level and severity of skin condition predicts outcome of psychotherapy in atopic dermatitis patients. <i>Int J Dermatol</i>, <i>40</i>, 632-636. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11737422">http://www.ncbi.nlm.nih.gov/pubmed/11737422</a></p>	<b>Atopic dermatitis (anxiety)</b>	<b>Brief Dynamic Psychotherapy</b> (Malan, 1992)	<b>Control (TAU?)</b>
109.	<p><b>Lipsitz, J. D., Gur, M., Vermes, D., Petkova, E., Cheng, J., Miller, N., ... Fyer, A. J. (2008).</b> A randomized trial of interpersonal therapy</p>	<b>Social anxiety disorder (SAD)</b>	<b>Psychodynamic Supportive Therapy</b> (Lipsitz et al., 2006)	<b>Interpersonal Therapy</b> (Klerman et al., 1984; Weissman et al., 2000)

	versus supportive therapy for social anxiety disorder. <i>Depression and Anxiety</i> , 25(6), 542–553. <a href="http://doi.wiley.com/10.1002/da.20364">http://doi.wiley.com/10.1002/da.20364</a>			
110.	<b>Lock, J., Grange, D. Le, &amp; Agras, W. (2010).</b> Randomized clinical trial comparing family-based treatment with adolescent-focused individual therapy for adolescents with anorexia nervosa. <i>Archives of General Psychiatry</i> , 67(10), 1025–1032. <a href="http://archpsyc.ama-assn.org/cgi/reprint/67/10/1025.pdf">http://archpsyc.ama-assn.org/cgi/reprint/67/10/1025.pdf</a>	<b>Anorexia</b>	<b>Adolescent Focused Therapy (AFT, Robin et al.)</b>	<b>Treatments Family-Based Treatment (FBT; Locke et al., 2001)</b>
111.	<b>Lorentzen, S., Ruud, T., Fjeldstad, A., &amp; Høglend, P. (2013).</b> Comparison of short- and long-term dynamic group psychotherapy: randomised clinical trial. <i>The British journal of psychiatry : the journal of mental science</i> . <a href="http://www.ncbi.nlm.nih.gov/pubmed/24029539">http://www.ncbi.nlm.nih.gov/pubmed/24029539</a>  <b>Lorentzen, S., Ruud, T., Fjeldstad, A., &amp; Høglend, P. a. (2014).</b> Personality disorder moderates outcome in short- and long-term group analytic psychotherapy: A randomized clinical trial. <i>The British Journal of Clinical Psychology / the British Psychological Society</i> , 1–18. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25178520">http://www.ncbi.nlm.nih.gov/pubmed/25178520</a>  <b>Fjeldstad, A., Høglend, P., &amp; Lorentzen, S. (2015).</b> Patterns of Change in Interpersonal Problems During and After Short-term and Long-term Psychodynamic Group Therapy: A Randomized Clinical Trial. <i>Psychotherapy Research</i> , 3307(November), 1–12. <a href="http://www.tandfonline.com/doi/full/10.1080/10503307.2015.1102357">http://www.tandfonline.com/doi/full/10.1080/10503307.2015.1102357</a>	<b>Anxiety and personality disorders</b>	<b>Long-Term Psychodynamic Group Therapy (Lorentzen, 2013)</b>	<b>Short-Term Psychodynamic Group Therapy (Lorentzen, 2013)</b>
112.	<b>Ludwig, G., Krenz, S., Zdrojewski, C., Bot, M., Rousselle, I., Stagno, D., ... Stiefel, F. (2013).</b> Psychodynamic interventions in cancer care I: psychometric results of a randomized controlled trial. <i>Psycho-oncology</i> , 23, 65-74. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23983096">http://www.ncbi.nlm.nih.gov/pubmed/23983096</a>	<b>Mixed depression and anxiety (in cancer patients)</b>	<b>Brief Dynamic Psychotherapy (Mann, 1973; Davanloo, 1994)</b>	<b>Waitlist</b>
113.	<b>Maina, G., Forner, F., &amp; Bogetto, F. (2005).</b> Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. <i>Psychother Psychosom</i> , 74, 43-50. <a href="http://www.ncbi.nlm.nih.gov/pubmed/15627856">http://www.ncbi.nlm.nih.gov/pubmed/15627856</a>	<b>Depression (mild to moderate)</b>	<b>Brief Dynamic Psychotherapy (BDP; Malan, 1979)</b>	<b>1. Brief Supportive Psychotherapy (BSP; Novalis et al., 1993)</b>  <b>2. Waitlist</b>
114.	<b>Maina, G., Rosso, G., Crespi, C., &amp; Bogetto, F. (2007).</b> Combined brief dynamic therapy and pharmacotherapy in the treatment of major depressive disorder: a pilot study. <i>Psychotherapy and Psychosomatics</i> , 76(5), 298–305. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17700050">http://www.ncbi.nlm.nih.gov/pubmed/17700050</a>  <b>Maina, G., Rosso, G., &amp; Bogetto, F. (2009).</b> Brief dynamic therapy combined with pharmacotherapy in the treatment of major depressive disorder: long-term results. <i>Journal of Affective Disorders</i> , 114(1-3), 200–7.	<b>Depression</b>	<b>Brief Dynamic Psychotherapy (BDP; Malan, 1963; 1976) + medication</b>	<b>Brief Supportive Psychotherapy (BSP; Novalis et al., 1993) + medication</b>

	<a href="http://www.ncbi.nlm.nih.gov/pubmed/18728001">http://www.ncbi.nlm.nih.gov/pubmed/18728001</a>			
115.	<b>Maina, G., Rosso, G., Rigardetto, S., Chiadò Piat, S., &amp; Bogetto, F. (2010).</b> No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. <i>Psychotherapy and Psychosomatics</i> , 79(5), 295–302. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20616624">http://www.ncbi.nlm.nih.gov/pubmed/20616624</a>	<b>OCD + depression</b>	<b>Brief Dynamic Psychotherapy (BDP; Malan, 1979) + medication</b>	<b>Medication only</b>
116.	<b>Marmar, C. R., Horowitz, M. J., Weiss, D. S., Wilner, N. R., &amp; Kaltreider, N. B. (1988).</b> A controlled trial of brief psychotherapy and mutual-help group treatment of conjugal bereavement. <i>Am J Psychiatry</i> , 145, 203-209. <a href="http://ps.psychiatryonline.org/data/Journals/AJP/3483/203.pdf">http://ps.psychiatryonline.org/data/Journals/AJP/3483/203.pdf</a>	<b>Bereavement</b>	<b>Brief Dynamic Psychotherapy (BDP; Malan, Sifneos; Mann)</b>	<b>Mutual Self-Help Groups</b>
117.	<b>Martini, B., Rosso, G., Chiodelli, D. F., De Cori, D., &amp; Maina, G. (2011).</b> Brief dynamic therapy combined with pharmacotherapy in the treatment of panic disorder with concurrent depressive symptoms. <i>Clinical Neuropsychiatry</i> , 8(3), 204–21. <a href="http://www.clinicalneuropsychiatry.org/pdf/04_martini.pdf">http://www.clinicalneuropsychiatry.org/pdf/04_martini.pdf</a>	<b>Panic disorder + depression</b>	<b>Brief Dynamic Psychotherapy (BDP; Malan, 1963) + medication</b>	<b>Brief Supportive Psychotherapy (BSP) + medication</b>
118.	<b>McCallum, M., &amp; Piper, W. E. (1990).</b> A controlled study of effectiveness and patient suitability for short-term group psychotherapy. <i>Int J Group Psychother</i> , 40, 431-452. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2228349">http://www.ncbi.nlm.nih.gov/pubmed/2228349</a>	<b>Mixed sample</b>	<b>Short-term psychoanalytically oriented group psychotherapy (STG; manual mentioned but no specific reference)</b>	<b>Waitlist</b>
119.	<b>McLean, P., &amp; Hakstian, A. (1979).</b> Clinical depression: comparative efficacy of outpatient treatments. <i>J Consult Clin Psychol</i> , 47, 818-836. <a href="http://www.ncbi.nlm.nih.gov/pubmed/389965">http://www.ncbi.nlm.nih.gov/pubmed/389965</a>	<b>Depression</b>	<b>Short-Term Psychotherapy (Marmor, 1975; Wolberg, 1967)</b>	<b>1. Relaxation therapy (no reference)</b> <b>2. Behavior therapy (McLean, 1976)</b> <b>3. Drug therapy</b>
120.	<b>McMain, S. F., Links, P. S., Gnam, W. H., Guimond, T., Cardish, R. J., &amp; Korman, L., et al. (2009).</b> A randomized trial of dialectical behavior therapy versus general psychiatric management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1365-1374. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19755574">http://www.ncbi.nlm.nih.gov/pubmed/19755574</a>  <b>McMain, S., Guimond, T., &amp; Streiner, D. (2012).</b> Dialectical Behavior Therapy Compared With General Psychiatric Management for Borderline Personality Disorder: Clinical Outcomes and Functioning Over a 2-Year Follow-Up. <i>American Journal of Psychiatry</i> , 650–661. <a href="http://www.neuro.psychiatryonline.org/article.aspx?articleid=1148702&amp;RelatedWidgetArticles=true">http://www.neuro.psychiatryonline.org/article.aspx?articleid=1148702&amp;RelatedWidgetArticles=true</a>	<b>Borderline personality disorder</b>	<b>General psychiatric management, dynamically informed therapy (Gunderson, 1984)</b>	<b>Dialectical Behavior Therapy (DBT; Linehan et al., 1991)</b>
121.	<b>Meyer, A. E., &amp; Bolz, W. (1981).</b> The Hamburg short psychotherapy comparison experiment. Foreword and introduction. <i>Psychother Psychosom</i> , 35, 81-84. <a href="http://www.karger.com/doi/10.1159/000287">http://www.karger.com/doi/10.1159/000287</a>	<b>Mixed sample</b>	<b>Psychodynamic conflict-centered time-limited psychotherapy (no reference)</b>	<b>Waitlist</b>

	<a href="#">484</a>			
122.	<b>Miller, L. C., Barrett, C. L., Hampe, E., &amp; Noble, H. (1972).</b> Comparison of reciprocal inhibition, psychotherapy, and waiting list control for phobic children. <i>Journal of Abnormal Psychology, 79</i> (3), 269–279. <a href="http://psycnet.apa.org/journals/abn/79/3/269">http://psycnet.apa.org/journals/abn/79/3/269</a>	<b>Phobia (children)</b>	<b>Play psychotherapy</b> (Lippman, 1956)	<b>Reciprocal inhibition</b>
123.	<b>Milrod, B., Chambless, D. L., Gallop, R., Busch, F. N., Schwalberg, M., McCarthy, K. S., ... Barber, J. P. (2015).</b> Psychotherapies for Panic Disorder: A Tale of Two Sites. <i>The Journal of Clinical Psychiatry, 100</i> 65, 1–9. <a href="http://www.psychiatrist.com/jcp/article/pages/2015/aheadofprint/14m09507.aspx">http://www.psychiatrist.com/jcp/article/pages/2015/aheadofprint/14m09507.aspx</a>	<b>Panic disorder</b>	<b>Panic-Focused Psychodynamic Psychotherapy (PFPP;</b> Milrod et al, 2001)	<b>1. CBT (Craske, 2000)</b> <b>2. Applied relaxation (Öst &amp; Westling, 1995)</b>
124.	<b>Milrod, B., Leon, A. C., Busch, F., Rudden, M., Schwalberg, M., &amp; Clarkin, J., et al. (2007).</b> A randomized controlled clinical trial of psychoanalytic psychotherapy for panic disorder. <i>Am J Psychiatry, 164</i> , 265-272. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17267789">http://www.ncbi.nlm.nih.gov/pubmed/17267789</a>	<b>Panic disorder</b>	<b>Panic-Focused Psychodynamic Psychotherapy (PFPP;</b> Milrod et al, 1997)	<b>Applied relaxation (Öst &amp; Westling, 1995)</b>
125.	<b>Monsen, K., &amp; Monsen, J. T. (2000).</b> Chronic Pain and Psychodynamic Body Therapy: A Controlled Outcome Study. <i>Psychotherapy: Theory, Research, Practice, Training, 37</i> (3), 257–269. <a href="http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&amp;id=2000-02713-007">http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&amp;id=2000-02713-007</a>	<b>Chronic pain</b>	<b>Psychodynamic body therapy (Monsen &amp; Monsen, 1999)</b>	<b>TAU</b>
126.	<b>Montag, C., Haase, L., Seidel, D., Bayerl, M., Gallinat, J., Herrmann, U., &amp; Dannecker, K. (2014).</b> A Pilot RCT of Psychodynamic Group Art Therapy for Patients in Acute Psychotic Episodes: Feasibility, Impact on Symptoms and Mentalising Capacity. <i>PLoS One, 9</i> (11), e112348. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4231093&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4231093&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Schizophrenia</b>	<b>Psychodynamic art therapy (Dannecker, 2010)</b>	<b>TAU</b>
127.	<b>Munroe-Blum, H., &amp; Marziali, E. (1995).</b> A controlled trial of short-term group treatment for borderline personality disorder. <i>J Personal Disord, 9</i> , 190-198. <a href="http://psycnet.apa.org/psycinfo/1996-17928-001">http://psycnet.apa.org/psycinfo/1996-17928-001</a>	<b>Borderline personality disorder</b>	<b>Time-limited group therapy (Marziali &amp; Munroe-Blum, 1995)</b>	<b>Individual psychodynamic psychotherapy (Kernberg, 1975)</b>
128.	<b>Muran, J. C., Safran, J. D., Samstag, L. W., &amp; Winston, A. (2005).</b> Evaluating an alliance-focused treatment for personality disorders. <i>Psychotherapy: Theory, Research, Practice, Training, 42</i> (4), 532–545. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.4.532">http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.4.532</a>  Safran, J. D., Muran, J. C., Samstag, L. W., & Winston, A. (2005). Evaluating alliance-focused intervention for potential treatment failures: A feasibility study and descriptive analysis. <i>Psychotherapy: Theory, Research, Practice, Training, 42</i> (4), 512–531. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.4.512">http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.4.512</a>	<b>Personality disorders (mixed; mainly cluster C)</b>	<b>Brief Relational Therapy (BRT; Safran &amp; Muran, 2000)</b>	<b>1. Short-Term Dynamic Psychotherapy (STDP: Pollock, Flegenheimer, &amp; Kaufman, 1992)</b> <b>2. Cognitive– behavioral therapy (CBT; Turner &amp; Muran, 1992; Beck, Freeman, &amp; Associates, 1990)</b>
129.	<b>Nader-Mohammadi Moghadam, M. Atef-Vahid, M. K., Asgharnejad-Farid, A. A.,</b>	<b>Social anxiety disorder (SAD)</b>	<b>Affect Phobia Therapy (APT; McCullough et al.,</b>	<b>1. Sertraline</b>

	Shabani, A., & Lavasni, F. (2015). Effectiveness of Short-term Dynamic Psychotherapy versus Sertraline in Treatment of Social Phobia. <i>Iran Journal of Psychiatry</i> , 9(2), 2–6. <a href="http://www.ncbi.nlm.nih.gov/pubmed/26288643">http://www.ncbi.nlm.nih.gov/pubmed/26288643</a>		2003)	<b>2. Waitlist</b>
130.	<b>Nijdam</b> , M. J., Gersons, B. P. R., Reitsma, J. B., de Jongh, A., & Olff, M. (2012). Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: randomised controlled trial. <i>The British Journal of Psychiatry: The Journal of Mental Science</i> , 200(3), 224–31. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22322458">http://www.ncbi.nlm.nih.gov/pubmed/22322458</a>	<b>PTSD</b>	<b>Brief eclectic psychotherapy (BEP;</b> Gersons et al, 2004; Manual includes both CBT and PDT components)	<b>Eye movement and reprocessing (EMDR;</b> De Jongh et al, 2004)
131.	<b>Ojehagen</b> , A., Berglund, M., Appel, C. P., Andersson, K., Nilsson, B., & Skjaeris, A., et al. (1992). A randomized study of long-term outpatient treatment in alcoholics. Psychiatric treatment versus multimodal behavioural therapy, during 1 versus 2 years of treatment. <i>Alcohol Alcohol</i> , 27, 649-658. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1292438">http://www.ncbi.nlm.nih.gov/pubmed/1292438</a>	<b>Alcohol dependence</b>	<b>“Psychiatric treatment” based on psychodynamic principles</b> (Luborsky, 1984)	<b>Multimodal behavior therapy</b> (Lazarus, 1981)
132.	<b>Pierloot</b> , R., & Vinck, J. (1978). Differential outcome of short-term dynamic psychotherapy and systematic desensitizations in the treatment of anxious outpatients: A preliminary report. <i>Psychologica Belgica</i> , 18, 87-98. <a href="http://psycnet.apa.org/psycinfo/1980-32873-001">http://psycnet.apa.org/psycinfo/1980-32873-001</a>	<b>Anxiety</b>	<b>Short-Term Dynamic Psychotherapy</b> (Malan, 1963)	<b>Systematic desensitization</b> (Wolpe, 1969)
133.	<b>Piper</b> , W. E., Azim, H. F., McCallum, M., & Joyce, A. S. (1990). Patient suitability and outcome in short-term individual psychotherapy. <i>J Consult Clin Psychol</i> , 58, 475-481. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2212185">http://www.ncbi.nlm.nih.gov/pubmed/2212185</a>	<b>Mixed sample</b>	<b>STDP</b> (Malan, 1976; Strupp & Binder, 1984)	<b>Waitlist</b>
134.	<b>Piper</b> , W. E., Debbane, E. G., Bienvenu, J. P., & Garant, J. (1984). A comparative study of four forms of psychotherapy. <i>J Consult Clin Psychol</i> , 52, 268-279. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6425378">http://www.ncbi.nlm.nih.gov/pubmed/6425378</a>	<b>Mixed sample</b>	<b>Short-term individual (STI;</b> Malan, 1976)	<b>1. Short-term group (STG)</b> <b>2. Long-term individual (LTI)</b> <b>3. Long-term group (LTG)</b>  Non manuals used
135.	<b>Piper</b> , W. E., Joyce, a S., McCallum, M., & Azim, H. F. (1998). Interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i> , 66(3), 558–67. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10224738">http://www.ncbi.nlm.nih.gov/pubmed/10224738</a>  <b>Piper</b> , W. E., McCallum, M., Joyce, a S., Azim, H. F., & Ogrodniczuk, J. S. (1999). Follow-up findings for interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i> , 67(2), 267–73. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10224738">http://www.ncbi.nlm.nih.gov/pubmed/10224738</a>	<b>Mixed sample</b>	<b>Interpretive psychotherapy</b> (Piper, Azim, Joyce, & McCallum, 1993)	<b>Supportive psychotherapy</b> (Piper, Azim, Joyce, & McCallum, 1993)
136.	<b>Piper</b> , W. E., McCallum, M., Joyce, A. S., Rosie,	<b>Complicated grief</b>	<b>Interpretive group</b>	<b>Supportive group therapy</b>

	<p>J. S., &amp; Ogrodniczuk, J. S. (2001). Patient personality and time-limited group psychotherapy for complicated grief. <i>Int J Group Psychother</i>, 51, 525-552. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11582899">http://www.ncbi.nlm.nih.gov/pubmed/11582899</a></p> <p>Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., Weideman, R., &amp; Rosie, J. S. (2007). Group composition and group therapy for complicated grief. <i>Journal of Consulting and Clinical Psychology</i>, 75(1), 116–25. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17295570">http://www.ncbi.nlm.nih.gov/pubmed/17295570</a></p> <p>Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., &amp; Weideman, R. (2009). Follow-up outcome in short-term group therapy for complicated grief. <i>Group Dynamics: Theory, Research, and Practice</i>, 13(1), 46–58. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0013830">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0013830</a></p>		therapy (no specific reference)	(no specific reference)
137.	<p><b>Poulsen, A.</b> (1991). Psychodynamic, time-limited group therapy in rheumatic disease--a controlled study with special reference to alexithymia. <i>Psychother Psychosom</i>, 56, 12-23. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1891575">http://www.ncbi.nlm.nih.gov/pubmed/1891575</a></p>	<b>Rheumatic disease</b>	<b>Time-limited group therapy</b> (Foulkes, 1975)	<b>Waitlist</b>
138.	<p><b>Poulsen, S., Lunn, S., Daniel, S. I. F., Folke, S., Mathiesen, B. B., Katznelson, H., &amp; Fairburn, C. G.</b> (2013). A Randomized Controlled Trial of Psychoanalytic Psychotherapy or Cognitive-Behavioral Therapy for Bulimia Nervosa. <i>The American journal of psychiatry</i>, 1–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24275909">http://www.ncbi.nlm.nih.gov/pubmed/24275909</a></p>	<b>Bulimia</b>	<b>Psychoanalytic Psychotherapy for bulimia nervosa</b> (Lunn & Poulsen, 2012)	<b>CBT</b> (Fairburn, 2008)
139.	<p><b>Reneses, B., Galián, M., Serrano, R., Figuera, D., Fernandez Del Moral, A., López-Ibor, J. J., ... Trujillo, M.</b> (2013). A new Time Limited Psychotherapy for BPD: Preliminary Results of a Randomized and Controlled Trial. <i>Actas españolas de psiquiatría</i>, 41(3), 139–48. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23803797">http://www.ncbi.nlm.nih.gov/pubmed/23803797</a></p>	<b>Borderline personality disorder</b>	<b>Representation focused Psychotherapy (PRFP;</b> (Trujillo, 2001; 2010)	<b>TAU</b>
140.	<p><b>Ridsdale, L., Godfrey, E., Chalder, T., Seed, P., King, M., Wallace, P., &amp; Wessely, S.</b> (2001). Chronic fatigue in general practice: is counselling as good as cognitive behaviour therapy? A UK randomised trial. <i>The British Journal of General Practice : The Journal of the Royal College of General Practitioners</i>, 51(462), 19–24. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1313894&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1313894&amp;tool=pmcentrez&amp;rendertype=abstract</a></p>	<b>Chronic fatigue</b>	<b>Counselling</b>  “...all counsellors used a psychodynamic approach and all cognitive behaviour therapists used CBT” (p. 20)	<b>CBT</b>
141.	<p><b>Robin, A L., Siegel, P. T., &amp; Moye, a.</b> (1995). Family versus individual therapy for anorexia: impact on family conflict. <i>The International Journal of Eating Disorders</i>, 17(4), 313–22. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7620470">http://www.ncbi.nlm.nih.gov/pubmed/7620470</a></p> <p>Robin, A. L., Siegel, P. T., Moye, A. W., Gilroy, M., Dennis, A. B., &amp; Sikand, A., et al. (1999). A controlled comparison of family versus individual therapy for adolescents with</p>	<b>Anorexia</b>	<b>Ego-oriented individual therapy (EOIT)</b> (Robin, Siegel, Moye, Koepke, & Tice, 1994)	<b>Behavioral family systems therapy (BFST;</b> Robin et al., 1994, 1996)

	<p>anorexia nervosa. <i>J Am Acad Child Adolesc Psychiatry</i>, 38, 1482-1489.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/10596247">http://www.ncbi.nlm.nih.gov/pubmed/10596247</a></p>			
142.	<p><b>Rosenbaum, B., Harder, S., Knudsen, P., Køster, A., Lindhardt, A., Lajer, M., ... Winther, G.</b> (2012). Supportive Psychodynamic Psychotherapy versus Treatment as Usual for First-Episode Psychosis: Two-Year Outcome. <i>Psychiatry</i>, 75(4), 331–41.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/23244011">http://www.ncbi.nlm.nih.gov/pubmed/23244011</a></p> <p>Harder, S., Koester, A., Valbak, K., &amp; Rosenbaum, B. (2014). Five-year follow-up of supportive psychodynamic psychotherapy in first-episode psychosis: long-term outcome in social functioning. <i>Psychiatry</i>, 77(2), 155–68.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/24865198">http://www.ncbi.nlm.nih.gov/pubmed/24865198</a></p>	<b>Schizophrenia</b>	<b>Supportive Psychodynamic Psychotherapy (SPP;</b> Rosenbaum, 2009)	<b>TAU</b>
143.	<p><b>Rosser, R., Denford, J., Heslop, A., Kinston, W., Macklin, D., &amp; Minty, K., et al.</b> (1983). Breathlessness and psychiatric morbidity in chronic bronchitis and emphysema: a study of psychotherapeutic management. <i>Psychol Med</i>, 13, 93-110.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/6844471">http://www.ncbi.nlm.nih.gov/pubmed/6844471</a></p>	<b>Breathlessness and psychiatric morbidity (in patients with chronic bronchitis and emphysema)</b>	<b>Analytical psychotherapy</b> (No specific reference)	<p><b>1. No treatment</b></p> <p><b>2. Nurse support</b></p>
144.	<p><b>Rosso, G., Martini, B., &amp; Maina, G.</b> (2012). Brief dynamic therapy and depression severity: A single-blind, randomized study. <i>Journal of affective disorders</i>, 1–6.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/23174499">http://www.ncbi.nlm.nih.gov/pubmed/23174499</a></p>	<b>Depression</b>	<b>Brief Dynamic Psychotherapy (BDP;</b> Malan, 1976)	<b>Brief Supportive Psychotherapy (BSP;</b> Novalis et al., 1993)
145.	<p><b>Rossouw, T. I., &amp; Fonagy, P.</b> (2012). Mentalization-based treatment for self-harm in adolescents: a randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i>, 51(12), 1304–1313.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/23200287">http://www.ncbi.nlm.nih.gov/pubmed/23200287</a></p>	<b>Self-harm (adolescents)</b>	<b>Mentalization-based treatment for adolescents (MBT-A;</b> Rossouw, unpublished)	<b>TAU</b>
146.	<p><b>Salminen, J. K., Karlsson, H., Hietala, J., Kajander, J., Aalto, S., &amp; Markkula, J., et al.</b> (2008). Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. <i>Psychother Psychosom</i>, 77, 351-357.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/18701831">http://www.ncbi.nlm.nih.gov/pubmed/18701831</a></p>	<b>Depression</b>	<b>Short-term psychodynamic psychotherapy (STDP;</b> Malan, 1976; Mann, 1973)	<b>Fluoxetine</b>
147.	<p><b>Salomonsson, B., &amp; Sandell, R.</b> (2011). A randomized controlled trial of mother–infant psychoanalytic treatment: I. Outcomes on self-report questionnaires and external ratings. <i>Infant Mental Health Journal</i>, 32(2), 207–231.  <a href="http://onlinelibrary.wiley.com/doi/10.1002/imhj.20291/full">http://onlinelibrary.wiley.com/doi/10.1002/imhj.20291/full</a></p>	<b>Mothers who worry about child</b>	<b>Mother–infant psychoanalytic treatment (MIP;</b> Norman, 2001; 2004)	<b>Child Health Center Care (TAU)</b>
148.	<p><b>Salzer, S., Cropp, C., Jaeger, U., Masuhr, O., &amp; Streeck-Fischer, a.</b> (2013). Psychodynamic therapy for adolescents suffering from co-morbid disorders of conduct and emotions in an in-patient setting: a randomized controlled trial. <i>Psychological medicine</i>, 1–10.</p>	<b>Mixed sample of adolescents</b>	<b>Psychoanalytic-interactive method (PiM;</b> Streeck & Leichsenring, 2009; Leichsenring et al., 2010)	<b>Waitlist/TAU</b>

	<a href="http://www.ncbi.nlm.nih.gov/pubmed/24229481">http://www.ncbi.nlm.nih.gov/pubmed/24229481</a>			
149.	<b>Sanchez, V. C., Lewinsohn, P. M., &amp; Larson, D. W. (1980).</b> Assertion training: effectiveness in the treatment of depression. <i>J Clin Psychol</i> , 36, 526-529. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7372826">http://www.ncbi.nlm.nih.gov/pubmed/7372826</a>	<b>Depression</b>	<b>"Traditional" Group Therapy</b> (No specific reference)	<b>Assertion training</b>
150.	<b>Sandahl, C., Herlitz, K., Ahlin, G., &amp; Rönnerberg, S. (1998).</b> Time-limited group psychotherapy for moderately alcohol dependent patients: A randomized controlled clinical trial. <i>Psychotherapy Research</i> , 8, 361-378. <a href="http://www.tandfonline.com/doi/abs/10.1080/10503309812331332467">http://www.tandfonline.com/doi/abs/10.1080/10503309812331332467</a>	<b>Alcohol dependence</b>	<b>Time-limited group psychotherapy</b> (Foulkes, 1964)	<b>Cognitive-behavioural group therapy</b> (Annis, 1986; Marlatt & Gordon, 1985; Sandahl and Rönnerberg, 1990)
151.	<b>Sandahl, C., Lundberg, U., Lindgren, A., Ryländer, G., Herlofson, J., Nygren, A., &amp; Asberg, M. (2011).</b> Two forms of group therapy and individual treatment of work-related depression: a one-year follow-up study. <i>International journal of group psychotherapy</i> , 61(4), 539–55. <a href="http://www.ncbi.nlm.nih.gov/pubmed/21985258">http://www.ncbi.nlm.nih.gov/pubmed/21985258</a>	<b>Depression</b> (work-related)	<b>Focused psychodynamic group therapy</b> (FGT; Sandahl & Lindgren, 2003)	<b>1. Cognitive Group Therapy</b> (no specific reference)  <b>2. Comparison condition</b>
152.	<b>Sattel, H., Lahmann, C., Gündel, H., Guthrie, E., Kruse, J., Noll-Hussong, M., ... Henningsen, P. (2011).</b> Brief psychodynamic interpersonal psychotherapy for patients with multisomatoform disorder: randomised controlled trial. <i>The British journal of psychiatry : the journal of mental science</i> , 60–67. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22075651">http://www.ncbi.nlm.nih.gov/pubmed/22075651</a>	<b>Multisomatoform disorder</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Guthrie, 2011; Henningsen, 2011)	<b>Enhanced medical care</b>
153.	<b>Saunders, D. G. (1996).</b> Feminist-cognitive-behavioral and process-psychodynamic treatments for men who batter: interaction of abuser traits and treatment models. <i>Violence Vict</i> , 11, 393-414. <a href="http://www.ncbi.nlm.nih.gov/pubmed/9210279">http://www.ncbi.nlm.nih.gov/pubmed/9210279</a>	<b>Violent men</b>	<b>Process-psychodynamic group therapy</b> (PPT; Browne, Saunders, & Staecker, in press)	<b>Feminist-group CBT</b> (FCBT; Browne, Saunders, & Staecker, in press)
154.	<b>Scheidt, C. E., Waller, E., Endorf, K., Schmidt, S., König, R., Zeeck, A., ... Lacour, M. (2012).</b> Is brief psychodynamic psychotherapy in primary fibromyalgia syndrome with concurrent depression an effective treatment? A randomized controlled trial. <i>General hospital psychiatry</i> . <a href="http://www.ncbi.nlm.nih.gov/pubmed/23218844">http://www.ncbi.nlm.nih.gov/pubmed/23218844</a>	<b>Fibromyalgia + depression</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Hobson, 1985; Goldberg et al., 1984; Guthrie, 2013)	<b>TAU</b>
155.	<b>Schnyder, U., &amp; Müller, J. (2011).</b> Brief Eclectic Psychotherapy for PTSD: A Randomized Controlled Trial. <i>The Journal of Clinical ...</i> , 564–566. <a href="http://europepmc.org/abstract/MED/21527127">http://europepmc.org/abstract/MED/21527127</a>	<b>PTSD</b>	<b>Brief eclectic psychotherapy</b> (BEP; Gersons et al, 2004; Manual includes both CBT and PDT components)	<b>Minimal attention control</b>
156.	<b>Shaffer, H. J., LaSalvia, T. A., &amp; Stein, J. P. (1997).</b> Comparing Hatha yoga with dynamic group psychotherapy for enhancing methadone maintenance treatment: a randomized clinical trial. <i>Altern Ther Health Med</i> , 3, 57-66. <a href="http://www.ncbi.nlm.nih.gov/pubmed/92107">http://www.ncbi.nlm.nih.gov/pubmed/92107</a>	<b>Opiate addiction</b>	<b>Dynamic group psychotherapy</b>	<b>Hatha yoga therapy</b>



	<a href="#">77</a>			
157.	<p><b>Shapiro, D. A., Barkham, M., Rees, A., Hardy, G. E., Reynolds, S., &amp; Startup, M., et al. (1994).</b> Effects of treatment duration and severity of depression on the effectiveness of cognitive-behavioral and psychodynamic-interpersonal psychotherapy. <i>J Consult Clin Psychol</i>, <i>62</i>, 522-534.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/8063978">http://www.ncbi.nlm.nih.gov/pubmed/8063978</a></p>	<b>Depression</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Hobson, 1985; Goldberg et al., 1984)	<b>Cognitive Therapy (CT;</b> Beck, Rush, Shaw, and Emery, 1979)
158.	<p><b>Shapiro, D., &amp; Firth, J. (1987).</b> Prescriptive v. Exploratory Psychotherapy Outcomes of the Sheffield Psychotherapy Project. <i>The British Journal of Psychiatry</i>, <i>151</i>, 790-799</p> <p><a href="http://bjp.publicaciones.saludcastillayleon.es/content/151/6/790.short">http://bjp.publicaciones.saludcastillayleon.es/content/151/6/790.short</a></p> <p>Shapiro, D. a, &amp; Firth-Cozens, J. (1990). Two-year follow-up of the Sheffield Psychotherapy Project. <i>The British journal of psychiatry</i>, <i>157</i>, 389–91.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/2245269">http://www.ncbi.nlm.nih.gov/pubmed/2245269</a></p>	<b>Deression and anxiety</b>	<b>Exploratory therapy</b> (no specific reference)	<b>Prescriptive therapy</b> (no specific reference)
159.	<p><b>Shaw, C. M., Margison, F. R., Guthrie, E. A., &amp; Tomenson, B. (2001).</b> Psychodynamic interpersonal therapy by inexperienced therapists in a naturalistic setting: A pilot study. <i>European Journal of Psychotherapy, Counselling, and Health</i>, <i>4</i>, 87–101.</p> <p><a href="http://www.tandfonline.com/doi/abs/10.1080/13642530110040091">http://www.tandfonline.com/doi/abs/10.1080/13642530110040091</a></p>	<b>Mixed sample</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Hobson, 1985)	<b>Waitlist</b>
160.	<p><b>Shefler, G., Dasberg, H., &amp; Ben-Shakhar, G. (1995).</b> A randomized controlled outcome and follow-up study of Mann's time-limited psychotherapy. <i>J Consult Clin Psychol</i>, <i>63</i>, 585-593.</p> <p><a href="http://psycnet.apa.org/journals/ccp/63/4/585">http://psycnet.apa.org/journals/ccp/63/4/585</a></p>	<b>Mixed sample</b>	<b>Time-Limited Psychotherapy</b> (TLP; Mann, 1973)	<b>Waitlist</b>
161.	<p><b>Simpson, S., Corney, R., Fitzgerald, P., &amp; Beecham, J. (2003).</b> A randomized controlled trial to evaluate the effectiveness and costeffectiveness of psychodynamic counselling for general practice patients with chronic depression. <i>Psychol Med</i>, <i>33</i>, 229-239.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/12622302">http://www.ncbi.nlm.nih.gov/pubmed/12622302</a></p>	<b>Depression (chronic)</b>	<b>Psychodynamic counselling</b> (Burton, 1998)	<b>General practice</b>
162.	<p><b>Sjodin, I., Svedlund, J., Ottosson, J. O., &amp; Dotevall, G. (1986).</b> Controlled study of psychotherapy in chronic peptic ulcer disease. <i>Psychosomatics</i>, <i>27</i>, 187-91, 195.</p> <p><a href="http://www.psychosomaticsjournal.com/article/S0033-3182(86)72711-4/abstract">http://www.psychosomaticsjournal.com/article/S0033-3182(86)72711-4/abstract</a></p>	<b>Peptic ulcer disease</b>	<b>Short-Term Dynamic Therapy</b> (Malan, 1976)	<b>Medication</b>
163.	<p><b>Sleed, M., Baradon, T., &amp; Fonagy, P. (2013).</b> New Beginnings for mothers and babies in prison: A cluster randomized controlled trial. <i>Attachment &amp; Human Development</i>, <i>15</i>(4), 349–67.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/23550526">http://www.ncbi.nlm.nih.gov/pubmed/23550526</a></p>	<b>High risk mothers (and their babies) in prison</b>	<b>Attachment-based intervention</b> (Baradon, 2006)	<b>Standard health and social care</b>
164.	<p><b>Sloane, R. B., Staples, F. R., Cristol, A. H., Yorkston, N. J., &amp; Whipple, K. (1975).</b> Short-term analytically oriented psychotherapy versus behavior therapy. <i>American Journal of</i></p>	<b>Anxiety or personality disorder</b>	<b>Short-Term Analytically Oriented Psychotherapy</b> (no specific reference)	<b>1. Waitlist</b> <b>2. Behavior Therapy</b> (no specific reference)

	<i>Psychiatry</i> , 132, 373-377. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1119588">http://www.ncbi.nlm.nih.gov/pubmed/1119588</a>			
165.	<b>Smyrniotis, K. X., &amp; Kirkby, R. J.</b> (1993). Long-term comparison of brief versus unlimited psychodynamic treatments with children and their parents. <i>J Consult Clin Psychol</i> , 61, 1020-1027. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8113479">http://www.ncbi.nlm.nih.gov/pubmed/8113479</a>	<b>Children with mixed problems</b>	<b>Time-limited brief psychotherapy</b> (Smyrniotis, Kirkby, Smyrniotis, & Picone, 1987)	<b>Time-unlimited psychotherapy</b> (Klein, 1975; Segal, 1978)
166.	<b>Snyder, D. K., Wills, R. M., &amp; Grady-Fletcher, a.</b> (1991). Long-term effectiveness of behavioral versus insight-oriented marital therapy: a 4-year follow-up study. <i>Journal of Consulting and Clinical Psychology</i> , 59(1), 138-41. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2002129">http://www.ncbi.nlm.nih.gov/pubmed/2002129</a>	<b>Couples with relational problems</b>	<b>Insight-Oriented Marital Therapy (IOMT;</b> no specific reference)	<b>Behavioral Marital Therapy (BMT;</b> no specific reference)
167.	<b>Sørensen, P., Birket-Smith, M., Wattar, U., Buemann, I., &amp; Salkovskis, P.</b> (2010). A randomized clinical trial of cognitive behavioural therapy versus short-term psychodynamic psychotherapy versus no intervention for patients with hypochondriasis. <i>Psychol Med</i> , 1-11. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20380779">http://www.ncbi.nlm.nih.gov/pubmed/20380779</a>	<b>Hypochondriasis</b>	<b>Short-Term Psychodynamic Psychotherapy</b> (No specific reference)	<b>1. Cognitive-behavioural Therapy (CBT;</b> Salkovskis, 1989; Warwick, 1989; Salkovskis et al. 2003) <b>2. Waitlist</b>
168.	<b>Stanton, A. H., Gunderson, J. G., Knapp, P. H., Frank, A. F., Vancelli, M. L., &amp; Schnitzer, R., et al.</b> (1984). Effects of psychotherapy in schizophrenia: I. Design and implementation of a controlled study. <i>Schizophr Bull</i> , 10, 520-563. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6151245">http://www.ncbi.nlm.nih.gov/pubmed/6151245</a>	<b>Schizophrenia (non-chronic)</b>	<b>Expressive, insight-oriented psychotherapy (EIO;</b> Stanton, in press)	<b>Reality-adaptive, supportive therapy (RAS;</b> May, 1968)
169.	<b>Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattes, H., ... Kruse, J.</b> (2016). Treating post-traumatic stress disorder by resource activation in Cambodia. <i>World Psychiatry</i> , 15(2), 182-183. <a href="http://doi.org/10.1002/wps.20312">http://doi.org/10.1002/wps.20312</a>	<b>PTSD</b>	<b>Resource-Oriented Trauma Therapy</b> (see article)	<b>Waitlist</b>
170.	<b>Svartberg, M., Seltzer, M. H., &amp; Stiles, T. C.</b> (1998). The effects of common and specific factors in short-term anxiety-provoking psychotherapy: a pilot process-outcome study. <i>J Nerv Ment Dis</i> , 186, 691-696. <a href="http://journals.lww.com/jonmd/Abstract/1998/11000/The_Effects_of_Common_and_Specific_Factors_in.5.aspx">http://journals.lww.com/jonmd/Abstract/1998/11000/The_Effects_of_Common_and_Specific_Factors_in.5.aspx</a>	<b>Mixed sample</b>	<b>Short-term anxiety-provoking psychotherapy (STAPP;</b> Sifneos, 1992)	<b>Non-directive Psychotherapy (NDP;</b> utvecklas specifikt för studien)
171.	<b>Svartberg, M., Stiles, T. C., &amp; Seltzer, M. H.</b> (2004). Randomized, controlled trial of the effectiveness of short-term dynamic psychotherapy and cognitive therapy for cluster C personality disorders. <i>Am J Psychiatry</i> , 161, 810-817. <a href="http://www.ncbi.nlm.nih.gov/pubmed/15121645">http://www.ncbi.nlm.nih.gov/pubmed/15121645</a>	<b>Personality disorder (Cluster C)</b>	<b>Short-Term Dynamic Psychotherapy (STDP;</b> McCullough Vaillant, 1997)	<b>Cognitive Therapy (CT;</b> Beck & Freeman, 1990)
172.	<b>Svedlund, J., Sjödin, I., Ottosson, J. O., &amp; Dotevall, G.</b> (1983). Controlled study of psychotherapy in irritable bowel syndrome. <i>Lancet</i> , 2, 589-592. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3240791&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3240791&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>IBS</b>	<b>Brief Dynamic Psychotherapy (Malan,</b> 1979)	<b>Treatment as usual (TAU)</b> (medical)

173.	<p><b>Szapocznik, J., Rio, A., Murray, E., Cohen, R., Scopetta, M., &amp; Rivas-Vazquez, A., et al. (1989).</b> Structural family versus psychodynamic child therapy for problematic Hispanic boys. <i>J.Consult.Clin.Psychol.</i>, 57, 571-578.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/2794177">http://www.ncbi.nlm.nih.gov/pubmed/2794177</a></p>	<b>Behavioral problems among boys</b>	<b>Individual psychodynamic child therapy (IPCT)</b> (Adams, 1974; Cooper & Wanerman, 1977)	<p><b>1. Structural Family Therapy (SFT;</b> Minuchin &amp; Fishman, 1981; Minuchin, Montalvo, Guerney, Rosman, &amp; Schuman, 1967)</p> <p><b>2. Recreational control condition</b></p>
174.	<p><b>Tasca, G., Ritchie, K., Conrad, G., Balfour, L., Gayton, J., &amp; Lybanon, V., et al. (2006).</b> Attachment scales predict outcome in a randomized controlled trial of two group therapies for binge eating disorder: An aptitude by treatment interaction. <i>Psychotherapy Research</i>, 16, 106-121.  <a href="http://www.tandfonline.com/doi/abs/10.1080/10503300500090928">http://www.tandfonline.com/doi/abs/10.1080/10503300500090928</a></p>	<b>Binge eating disorder</b>	<b>Group psychodynamic interpersonal psychotherapy</b> (Tasca et al., 2002; Strupp & Binder, 1984; Malan 1979)	<b>Group cognitive-behavior therapy (GCBT;</b> Wilfley et al., 1996)
175.	<p><b>Thompson, L. W., Gallagher, D., &amp; Breckenridge, J. S. (1987).</b> Comparative effectiveness of psychotherapies for depressed elders. <i>J Consult Clin Psychol</i>, 55, 385-390.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/3597953">http://www.ncbi.nlm.nih.gov/pubmed/3597953</a></p>	<b>Depression (among elderly)</b>	<b>Brief psychodynamic therapy</b> (Horowitz & Kaltreider, 1979)	<p><b>1. Cognitive Therapy (CT;</b> Beck, Rush, Shaw, &amp; Emery, 1979)</p> <p><b>2. Behavioral Therapy (BT;</b> Gallagher &amp; Thompson, 1981; Lewinsohn, 1974)</p> <p><b>3. Delayed-treatment control</b></p>
176.	<p><b>Thyme, K. E., Sundin, E. C., Stahlberg, G., Lindstrom, B., Eklof, H., &amp; Wiberg, B., et al. (2007).</b> The outcome of short-term psychodynamic art therapy compared to short-term psychodynamic verbal therapy for depressed women. <i>Psychoanalytic Psychotherapy</i>, 21, 250-264.  <a href="http://www.tandfonline.com/doi/abs/10.1080/02668730701535610">http://www.tandfonline.com/doi/abs/10.1080/02668730701535610</a></p>	<b>Depression (among women)</b>	<b>Short-Term Psychodynamic Verbal Therapy</b> (Mann, 1973)	<b>Art psychotherapy</b> (no reference)
177.	<p><b>Trowell, J., Joffe, I., Campbell, J., Clemente, C., Almqvist, F., &amp; Soininen, M., et al. (2007).</b> Childhood depression: a place for psychotherapy. An outcome study comparing individual psychodynamic psychotherapy and family therapy. <i>Eur Child Adolesc Psychiatry</i>, 16, 157-167.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/17200793">http://www.ncbi.nlm.nih.gov/pubmed/17200793</a></p> <p>Garoff, F. F., Heinonen, K., Pesonen, A.-K., &amp; Almqvist, F. (2012). Depressed youth: treatment outcome and changes in family functioning in individual and family therapy. <i>Journal of Family Therapy</i>, 34(1), 4–23.  <a href="http://doi.wiley.com/10.1111/j.1467-6427.2011.00541.x">http://doi.wiley.com/10.1111/j.1467-6427.2011.00541.x</a></p>	<b>Depression (children)</b>	<b>Focused Individual Psychotherapy</b> (Malan & Osmimo, 1992; Davanloo, 1978)	<b>Systems Integrative Family Therapy</b> (Byung-Hall, 1995; Will, 1985)
178.	<p><b>Trowell, J., Kolvin, I., Weeramanthri, T., Sadowski, H., Berelowitz, M., &amp; Glaser, D., et al. (2002).</b> Psychotherapy for sexually abused girls: psychopathological outcome findings and patterns of change. <i>Br J Psychiatry</i>, 180, 234-247.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/11872516">http://www.ncbi.nlm.nih.gov/pubmed/11872516</a></p>	<b>Trauma (sexually abused girls)</b>	<b>Individual therapy</b> (Unclear with no reference)	<b>Psychopedagogical /analytic group therapy</b>
179.	<p><b>Vinnars, B., Barber, J. P., Norén, K., Gallop, R., &amp; Weinryb, R. M. (2005).</b> Manualized</p>	<b>Personality disorder (mixed)</b>	<b>Supportive-Expressive Psychotherapy (SE;</b>	<b>Non-manualized PDT</b>

	<p>supportive-expressive psychotherapy versus nonmanualized community-delivered psychodynamic therapy for patients with personality disorders: bridging efficacy and effectiveness. <i>Am J Psychiatry</i>, 162, 1933-1940.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/16199841">http://www.ncbi.nlm.nih.gov/pubmed/16199841</a></p> <p>Vinnars, B., Gallop, R., &amp; Barber, J. P. (2009). Do Personality Problems Improve During Psychodynamic Supportive-Expressive Psychotherapy? Secondary Outcome Results From a Randomized Controlled Trial for Psychiatric Outpatients with Personality Disorders. <i>Psychotherapy</i>, 46, 362-375.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/20161588">http://www.ncbi.nlm.nih.gov/pubmed/20161588</a></p>		Luborsky, 1984)	
180.	<p><b>Vitriol, V. G., Ballesteros, S. T., Florenzano, R. U., Weil, K. P., &amp; Benadof, D. F. (2009).</b> Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. <i>Psychiatr Serv</i>, 60, 936-942.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/19564224">http://www.ncbi.nlm.nih.gov/pubmed/19564224</a></p>	<b>Depression (+ childhood trauma)</b>	<b>Brief psychodynamic psychotherapy</b> (Kudler et al, 2003; Davies et al, 1994)	<b>TAU</b>
181.	<p><b>Watzke, B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., &amp; Schulz, H. (2010).</b> Effectiveness of systematic treatment selection for psychodynamic and cognitive-behavioural therapy: randomised controlled trial in routine mental healthcare. <i>The British Journal of Psychiatry : The Journal of Mental Science</i>, 197(2), 96-105.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/20679260">http://www.ncbi.nlm.nih.gov/pubmed/20679260</a></p> <p>Watzke, B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., &amp; Schulz, H. (2012). Longer term outcome of cognitive-behavioural and psychodynamic psychotherapy in routine mental health care: Randomised controlled trial. <i>Behaviour research and therapy</i>, 50(9), 580-587.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/22750189">http://www.ncbi.nlm.nih.gov/pubmed/22750189</a></p>	<b>Mixed sample</b>	<b>Psychodynamic brief group therapy</b> (Heigl-Evers et al., 1997)	<b>Cognitive-behavioural group therapy</b> (Fiedler, 2001)
182.	<p><b>Wiborg, I. M., &amp; Dahl, A. A. (1996).</b> Does brief dynamic psychotherapy reduce the relapse rate of panic disorder? <i>Arch Gen Psychiatry</i>, 53, 689-694.  <a href="http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1996.01830080041008">http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1996.01830080041008</a></p>	<b>Panic disorder</b>	<b>Brief Dynamic Psychotherapy (BDP;</b> Davanloo, 1978; Malan 1976; 1979; Strupp & Binder, 1984)	<b>Clomipramin alone</b>
183.	<p><b>Wiltink, J., Dippel, A., Szczepanski, M., Thiede, R., Alt, C., &amp; Beutel, M. E., et al. (2007).</b> Long-term weight loss maintenance after inpatient psychotherapy of severely obese patients based on a randomized study: predictors and maintaining factors of health behavior. <i>J Psychosom Res</i>, 62, 691-698.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/17540227">http://www.ncbi.nlm.nih.gov/pubmed/17540227</a></p>	<b>Severe obesity</b>	<b>Psychodynamic inpatient treatment</b> (no manual but reference to: Levine & Luborsky, 1981)	<b>Behavioral treatment</b> (no specific reference)
184.	<p><b>Winston, A., Pollack, J., McCullough, L., Flegenheimer, W., Kestenbaum, R., &amp; Trujillo, M., et al. (1991).</b> Brief psychotherapy of personality disorders. <i>J Nerv Ment Dis</i>, 179,</p>	<b>Personality disorders</b> (mixed)	<b>Short-Term Dynamic Therapy</b> (STDP; Davanloo, 1980; Also reference to Mann, Malan, and Sifneos)	<b>Brief Adaptive Psychotherapy</b> (BAP; Pollack et al., 1991)

	<p>188-193.  <a href="http://psycnet.apa.org/psycinfo/1991-25198-001">http://psycnet.apa.org/psycinfo/1991-25198-001</a></p> <p>Winston, A, Laikin, M., Pollack, J., Samstag, L. W., McCullough, L., &amp; Muran, J. C. (1994). Short-term psychotherapy of personality disorders. <i>The American journal of psychiatry</i>, 151(2), 190-4.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/22217220">http://www.ncbi.nlm.nih.gov/pubmed/22217220</a></p>			
185.	<p><b>Woody</b>, G. E., Luborsky, L., McLellan, A. T., O'Brien, C. P., Beck, A. T., &amp; Blaine, J., et al. (1983). Psychotherapy for opiate addicts. Does it help? <i>Arch Gen Psychiatry</i>, 40, 639-645.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/6847332">http://www.ncbi.nlm.nih.gov/pubmed/6847332</a></p> <p>Woody, G. E., Luborsky, L., McLellan, A. T., &amp; O'Brien, C. P. (1990). Corrections and Revised analyses for Psychotherapy in Methadone Patients. <i>Arch Gen Psychiatry</i>, 47, 788-789.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/2378551">http://www.ncbi.nlm.nih.gov/pubmed/2378551</a></p>	<b>Opiate dependence</b>	<b>Supportive-Expressive Psychotherapy</b> (SE; Luborsky, 1977; unpublished manual)	<b>Cognitive-Behavior Therapy</b> (CBT; Beck, unpublished)  <b>Drug Counseling</b> (DC; Stockdale & Hargrove, unpublished material, 1977)
186.	<p><b>Woody</b>, G. E., McLellan, A. T., Luborsky, L., &amp; O'Brien, C. P. (1995). Psychotherapy in community methadone programs: A validation study. <i>American Journal of Psychiatry</i>, 152, 1302-1308.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/7653685">http://www.ncbi.nlm.nih.gov/pubmed/7653685</a></p>	<b>Opiate dependence</b>	<b>Supportive-Expressive Psychotherapy</b> (SE; Luborsky, 1984)	<b>Drug counseling</b>
187.	<p><b>Zipfel</b>, S., Wild, B., Groß, G., Friederich, H., Teufel, M., Schellberg, D., ... Zwaan, M. De. (2013). Articles Focal psychodynamic therapy, cognitive behaviour therapy, and optimised treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised controlled trial. <i>The Lancet</i>, 383, 127-137.  <a href="http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)61746-8/abstract">http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)61746-8/abstract</a></p>	<b>Anorexia</b>	<b>Focal psychodynamic therapy</b> (FPT; Schauenburg et al., 2009)	<b>1. Enhanced cognitive behaviour therapy</b> (CBT-E; Fairburn, 2008)  <b>2. Optimised TAU</b>